Journal of Rural Advancement
[ISSN 2347-2561 (P); 2583-6102 (E)]
Vol. 12, Issue 02, October 2024

Harnessing Plant–Microbe Interactions for Sustainable Agricultural Transformation in Rural India

Prateek Jain

Department of Biotechnology and Bioinformatics, North-Eastern Hill University, Shillong 793022, Meghalaya, India
Email: prateekjain20000@gmail.com
Received on: June 10, 2024 ; Accepted on: July 3, 2024

Abstract

Indian agriculture stands at a critical ecological and socioeconomic crossroads. While the Green Revolution enabled national food self-sufficiency, decades of intensive chemical fertilizer and pesticide application, monocropping, and groundwater depletion have precipitated severe soil organic carbon (SOC) loss, secondary salinization, declining factor productivity, and environmental degradation across rural agroecosystems. Concurrently, climate change has intensified drought, heatwaves, and erratic monsoons, disproportionately affecting smallholder and marginal farmers who cultivate over 86% of India's operational landholdings. Harnessing beneficial plant–microbe interactions—encompassing plant growth-promoting rhizobacteria (PGPR), arbuscular mycorrhizal fungi (AMF), nitrogen-fixing diazotrophs, endophytes, and biological control agents—offers an ecologically sound, economically viable pathway to restore soil health, enhance nutrient-use efficiency (NUE), confer abiotic and biotic stress resilience, and reduce agrochemical dependency. This review synthesizes the mechanistic foundations of rhizosphere and endosphere symbioses, evaluates region-specific translational opportunities across India's diverse agro-climatic zones (from the Indo-Gangetic Plains and rainfed drylands to the acidic soils of the North-Eastern Himalayas), and critically examines the biological, technological, regulatory, and socioeconomic bottlenecks hindering large-scale rural adoption. Finally, we outline a forward-looking roadmap integrating multi-omics-guided synthetic microbial communities (SynComs), nano-encapsulated and biofilm-based bioformulations, farmer producer organization (FPO)-led decentralized biomanufacturing, and supportive policy reforms to drive a second, evergreen agricultural transformation in rural India.

Keywords: Plant–microbe interactions; Biofertilizers; Plant growth-promoting rhizobacteria (PGPR); Arbuscular mycorrhizal fungi (AMF); Sustainable agriculture; Smallholder farmers; Rural India; Synthetic microbial communities (SynComs)

1. Introduction

Agriculture remains the socioeconomic backbone of rural India, supporting the livelihoods of nearly half of the nation's workforce and contributing approximately 18% to the gross domestic product (GDP). Following the Green Revolution of the late 1960s, the widespread adoption of high-yielding crop varieties, assured irrigation, and synthetic agrochemicals propelled India from chronic food deficits to a global agricultural powerhouse (Pingali, 2012). However, the long-term ecological externalities of this input-intensive paradigm have become increasingly unsustainable. Decades of imbalanced

nitrogen–phosphorus–potassium (NPK) fertilization—heavily skewed toward subsidized urea—have depleted soil organic matter, disrupted native soil microbial biodiversity, acidified or salinized arable soils, and caused widespread multi-nutrient deficiencies, particularly in zinc, iron, boron, and sulfur (Bijay-Singh & Craswell, 2021; Bhattacharyya et al., 2015). Furthermore, the partial factor productivity of synthetic fertilizers in major cropping systems such as the rice–wheat rotation of the Indo-Gangetic Plains has declined precipitously, forcing farmers to apply escalating doses of chemical inputs merely to maintain baseline yields, thereby eroding farm profitability and contaminating groundwater with nitrates and pesticide residues (Ladha et al., 2003; Abrol et al., 2017).

These soil degradation challenges are compounded by the structural realities of rural Indian farming and accelerating climate change. Approximately 86.2% of Indian farmers are smallholders or marginal cultivators owning less than two hectares of land, operating with limited capital reserves and high vulnerability to input price volatility and climatic shocks (Birthal et al., 2015). Nearly 51% of India's net sown area remains rainfed, exposing crops to recurrent droughts, terminal heat stress, soil salinity, and emerging pest and pathogen outbreaks (Venkateswarlu & Shanker, 2009). Simultaneously, heavy reliance on imported potassic and phosphatic raw materials places a massive fiscal burden on the national exchequer through fertilizer subsidies, underscoring the urgent strategic imperative for domestic, bio-based alternatives (Sapkota & Bijay-Singh, 2025).

In nature, plants do not function as autonomous entities; rather, they operate as integrated ecological units—termed holobionts—co-evolving with complex communities of bacteria, fungi, archaea, and protists colonizing the rhizosphere (root–soil interface), rhizoplane (root surface), endosphere (internal tissues), and phyllosphere (aerial surfaces) (Vandenkoornhuyse et al., 2015; Bulgarelli et al., 2013). Through sophisticated chemical signaling mediated by root exudates—comprising organic acids, sugars, amino acids, flavonoids, and strigolactones—plants actively recruit beneficial microbiota that reciprocate by mobilizing locked soil nutrients, fixing atmospheric nitrogen, synthesizing phytohormones, priming systemic immunity, and buffering abiotic stresses (Bais et al., 2006; Sasse et al., 2018). For rural India, translating these plant–microbe interactions into low-cost, locally adapted biofertilizers, biostimulants, and biopesticides represents a transformative opportunity to decouple agricultural productivity from ecological degradation (Gupta et al., 2021; Singh et al., 2016).

Despite decades of foundational microbiological research across Indian agricultural universities and national institutes, the field-level translation of microbial technologies remains uneven. Commercial biofertilizers currently substitute only a fraction of chemical fertilizer demand, constrained by inconsistent field efficacy across heterogeneous agro-climatic zones, short shelf life under tropical storage conditions, suboptimal carrier formulations, quality control deficits, and weak last-mile extension networks (Vessey, 2003; Herrmann & Lesueur, 2013; Kumar et al., 2021). This review provides a comprehensive, critical synthesis of the mechanisms governing beneficial plant–microbe symbioses, evaluates their agronomic potential across India's major rural agroecosystems, dissects the technical and institutional barriers to adoption, and proposes next-generation biotechnological and policy interventions to mainstream

microbiome-driven sustainable agriculture in rural India.

2. Mechanistic Foundations of Beneficial Plant–Microbe Interactions

Beneficial plant-associated microorganisms—broadly categorized into plant growth-promoting rhizobacteria (PGPR), nitrogen-fixing diazotrophs, arbuscular mycorrhizal fungi (AMF), plant growth-promoting fungi (PGPF), and endophytes—facilitate plant growth and health via interconnected direct and indirect physiological pathways (Glick, 2012; Bhattacharyya & Jha, 2012) (Table 1).




Figure-1. Ectomycorrhizal vs. endomycorrhizal root colonization.

As illustrated in the mycorrhizal colonization diagram above, endomycorrhizal fungi penetrate root cortical cells to form branching arbuscules and external hyphae that expand nutrient absorption beyond the root zone, complementing rhizosphere bacteria and soil-borne antagonists across the soil–root continuum.

2.1 Nutrient Acquisition and Biogeochemical Cycling

Nitrogen (N) and phosphorus (P) are the primary limiting macronutrients across Indian agricultural soils, yet conventional synthetic applications suffer from dismal recovery efficiencies—typically 30–40% for applied N and only 15–20% for applied P (Tilman et al., 2002). Biological nitrogen fixation (BNF) represents the most sustainable alternative to synthetic Haber–Bosch nitrogen. Symbiotic rhizobia (*Rhizobium*, *Bradyrhizobium*, *Mesorhizobium*, *Sinorhizobium*) establish specialized root nodules in leguminous crops (chickpea, pigeonpea, mungbean, blackgram, groundnut, soybean), wherein host-secreted flavonoids activate bacterial NodD regulators, triggering Nod factor synthesis and root hair curling (Oldroyd, 2013; Herridge et al., 2008). Inside oxygen-buffered symbiosomes, bacteroids express the nitrogenase complex to reduce atmospheric N_2 into bioavailable ammonia. In non-leguminous cereals (rice, wheat, maize, millets) and cash crops like sugarcane, associative and endophytic diazotrophs—including *Azospirillum brasilense*, *Azotobacter chroococcum*, *Herbaspirillum seropedicae*, and *Gluconacetobacter diazotrophicus*—colonize root surfaces and intercellular spaces, contributing substantial fixed nitrogen while minimizing leaching and

nitrous oxide (N_2O) emissions (Boddey et al., 2003; Santi et al., 2013).

Functional Category	Key Microbial Taxa	Primary Biochemical Mechanisms	Agronomic Outcomes in Indian Crops	Representative References
Symbiotic & Associative N_2 Fixers	*Rhizobium*, *Bradyrhizobium*, *Mesorhizobium*, *Azospirillum*, *Azotobacter*, *Gluconacetobacter*	Reduction of atmospheric NH_2 to NH_3 via oxygen-sensitive nitrogenase enzyme complex (*nif* genes)	Supplies 20–40 kg N/ha in cereals/sugarcane and up to 100–150 kg N/ha in pulses/groundnut	Herridge et al. (2008); Boddey et al. (2003); Santi et al. (2013)
Phosphate & Potassium Solubilizers	*Bacillus megaterium*, *Pseudomonas striata*, *Pantoea*, *Aspergillus niger*, *Penicillium*, *Frateuria aurantia*	Secretion of gluconic, citric, and oxalic acids; release of acid/alkaline phosphatases and phytases	Unlocks insoluble Ca-, Fe-, and Al-bound phosphorus in alkaline and acidic Indian soils	Richardson & Simpson (2011); Alori et al. (2017); Meena et al. (2014)
Mycorrhizal Symbionts	*Funneliformis mosseae*, *Rhizophagus irregularis*, *Gigaspora*, *Glomus* spp.	Extraradical hyphal network expansion; glomalin-related soil protein (GRSP) secretion; polyphosphate transport	Enhances P, Zn, and water uptake by 30–60%; stabilizes soil aggregates in drylands	Smith & Read (2008); Rillig (2004); Auge (2001)
Phytohormone & ACC Deaminase Producers	*Pseudomonas fluorescens*, *Bacillus subtilis*, *Enterobacter*, *Azospirillum brasilense*	Biosynthesis of IAA, gibberellins, and cytokinins; cleavage of ACC into α-ketobutyrate and NH_3	Stimulates lateral root proliferation; lowers stress-induced ethylene under drought and salinity	Glick et al. (2007); Spaepen et al. (2007)
Biological Control Agents (BCAs)	*Trichoderma harzianum*, *T. viride*, *Pseudomonas fluorescens*, *Bacillus amyloliquefaciens*, *Streptomyces*	Mycoparasitism (chitinases, β-1,3-glucanases), antibiosis (2,4-DAPG, phenazines, lipopeptides), ISR priming	Suppresses *Fusarium* wilt, *Rhizoctonia* root rot, *Pythium* damping-off, and bacterial blight	Harman et al. (2004); Haas & Défago (2005); Pieterse et al. (2014)

Phosphorus in Indian soils is predominantly immobilized as insoluble tricalcium phosphate in alkaline/calcareous soils (Indo-Gangetic Plains and Deccan Plateau) or bound to iron and aluminum oxides in acidic soils (North-Eastern and Eastern India). Phosphate-solubilizing microorganisms (PSMs)—including *Bacillus megaterium*, *Pseudomonas putida*, *Enterobacter*, Aspergillus, and Penicillium—solubilize inorganic phosphates by secreting low-molecular-weight organic acids (gluconic, 2-ketogluconic, citric, malic, and oxalic

acids) that chelate Ca2+, Fe3+, and Al3+ cations and lower rhizosphere pH (Rodríguez & Fraga, 1999; Alori et al., 2017). Concurrently, they mineralize organic phosphorus pools via extracellular phytases and acid/alkaline phosphor-monoesterases (Richardson & Simpson, 2011). Similarly, potassium-solubilizing bacteria (Frateuria aurantia, Bacillus mucilaginosus) weather silicate minerals such as mica and feldspar via acidolysis and capsular polysaccharide complexation, reducing India's dependence on imported muriate of potash (Meena et al., 2014).

Arbuscular mycorrhizal fungi (phylum Glomeromycota, e.g., *Rhizophagus irregularis*, *Funneliformis mosseae*) form obligate endosymbioses with over 80% of terrestrial crop species. Upon perceiving host-derived strigolactones, AMF hyphae penetrate root cortical cells to form highly branched arbuscules—specialized interfaces for bidirectional nutrient exchange—while extending a vast extraradical hyphal network far beyond the root phosphorus-depletion zone (Smith & Read, 2008; Parniske, 2008). Beyond phosphorus, AMF hyphae efficiently scavenge immobile micronutrients such as zinc (Zn) and copper (Cu), directly addressing widespread micronutrient malnutrition ("hidden hunger") in Indian staple grains (Lehmann et al., 2014; Singh & Prasanna, 2020). Furthermore, high-affinity microbial siderophores (catecholates, hydroxamates, and carboxylates) produced by *Pseudomonas* and *Bacillus* chelate insoluble ferric iron (Fe^{3+}) in calcareous soils, facilitating plant iron acquisition while starving phytopathogens of essential iron (Kloepper et al., 1980; Saha et al., 2016).

2.2 Phytohormonal Modulation and Root System Architecture

Beneficial microbes profoundly remodel plant root architecture—increasing lateral root density, root hair elongation, and total absorptive surface area—by modulating endogenous phytohormone homeostasis (Spaepen et al., 2007). Over 80% of rhizosphere bacteria synthesize indole-3-acetic acid (IAA) primarily via the tryptophan-dependent indole-3-pyruvate (IPyA) pathway; low-to-moderate bacterial IAA concentrations stimulate primary root elongation and lateral root primordia initiation, enhancing water and nutrient foraging in nutrient-poor soils (Spaepen et al., 2007; Patten & Glick, 2002). Additionally, PGPR and PGPF produce gibberellins (promoting stem elongation and seed germination) and cytokinins (stimulating cell division and delaying leaf senescence) (Arkhipova et al., 2005).

A pivotal mechanism under environmental stress is the enzymatic cleavage of 1-aminocyclopropane-1-carboxylate (ACC), the immediate biosynthetic precursor of ethylene in higher plants. Under drought, salinity, waterlogging, or pathogen attack, plants synthesize deleterious bursts of "stress ethylene" that inhibit root elongation and accelerate senescence. PGPR possessing the enzyme **ACC deaminase** (encoded by *acdS*, common in *Pseudomonas*, *Burkholderia*, *Variovorax*, and *Enterobacter*) sequester exuded ACC from plant roots and hydrolyze it into ammonia and α-ketobutyrate, thereby dampening ethylene accumulation and sustaining root growth under adverse field conditions (Glick et al., 2007; Glick, 2014).

2.3 Abiotic Stress Mitigation: Drought, Salinity, and Temperature Extremes

Climate-induced abiotic stresses severely constrain crop yields across rural India,

where ~6.73 million hectares are affected by salinity and sodicity and over two-thirds of agricultural land is prone to periodic drought (Sharma et al., 2016). Halotolerant and drought-adapted PGPR and AMF confer systemic abiotic stress tolerance through coordinated physiological interventions (Yang et al., 2009; Dodd & Pérez-Alfocea, 2012). First, microbes trigger osmotic adjustment by inducing plant accumulation of compatible solutes—proline, glycine betaine, trehalose, and soluble sugars—that stabilize proteins and cellular membranes without disrupting enzymatic activity (Bano & Fatima, 2009; Kaushal & Wani, 2016). Second, under saline conditions, beneficial rhizobacteria regulate host ion transporters (such as the high-affinity potassium transporter *HKT1* and salt overly sensitive *SOS1* antiporters), restricting toxic Na^+ influx into shoots while maintaining a favorable cytosolic $K^+\backslash Na^+$ ratio (Zhang et al., 2008; Bharti et al., 2016).

Third, microbial inoculation upregulates plant antioxidant defense enzymes—including superoxide dismutase (SOD), catalase (CAT), ascorbate peroxidase (APX), and glutathione reductase (GR)—scavenging cytotoxic reactive oxygen species (ROS) generated during drought and heat stress (Gururani et al., 2013). Fourth, bacterial exopolysaccharides (EPS) form hydrophilic biofilms around root surfaces, binding Na^+ ions, enhancing rhizosheath water retention, and improving soil macro-aggregation alongside AMF-secreted glomalin-related soil proteins (GRSP) (Sandhya et al., 2009; Rillig, 2004; Auge, 2001).

2.4 Biocontrol of Phytopathogens and Induced Systemic Resistance

Soil-borne fungal, bacterial, and nematode pathogens—including *Fusarium oxysporum*, *Rhizoctonia solani*, *Macrophomina phaseolina*, *Sclerotium rolfsii*, *Xanthomonas oryzae*, and root-knot nematodes (*Meloidogyne* spp.)—inflict annual crop losses of 15–25% in India. Biological control agents (BCAs) suppress disease through both direct antagonism and host immune priming (Compant et al., 2005; Köhl et al., 2019). *Trichoderma* species (*T. harzianum*, *T. viride*, *T. asperellum*), which account for the majority of commercial biofungicides in India, exhibit aggressive mycoparasitism: they sense pathogen-derived oligochitins, coil around host hyphae, form appressoria, and secrete cell wall-degrading enzymes (chitinases, β-1,3-glucanases, and proteases) alongside antifungal peptaibols, gliotoxin, and plant-derived bioactive compounds with complementary antibacterial and antifungal properties (Harman et al., 2004; Druzhinina et al., 2011; Tyśkiewicz et al., 2022; Sharma, 2024).

Fluorescent pseudomonads (*P. fluorescens*, *P. protegens*) and *Bacillus* species (*B. subtilis*, *B. amyloliquefaciens*, *B. velezensis*) produce broad-spectrum antimicrobial metabolites, including 2,4-diacetylphloroglucinol (2,4-DAPG), phenazines, pyrrolnitrin, pyoluteorin, hydrogen cyanide (HCN), and cyclic lipopeptides (surfactin, iturin, fengycin) that disrupt pathogen membrane integrity (Haas & Défago, 2005; Ongena & Jacques, 2008). Crucially, colonization by non-pathogenic PGPR and *Trichoderma* triggers Induced Systemic Resistance (ISR) throughout the entire plant. Unlike pathogen-induced systemic acquired resistance (SAR), which relies on salicylic acid (SA) and pathogenesis-related (PR) proteins, ISR is predominantly mediated by jasmonic acid (JA) and ethylene (ET) signaling cascades, priming the host for

rapid callose deposition, phenolic accumulation, and phytoalexin synthesis upon subsequent attack by foliar fungi, bacteria, or insect herbivores (Pieterse et al., 2014; Van Wees et al., 2008).

3. Opportunities Across India's Diverse Rural Agroecosystems

India's vast pedoclimatic heterogeneity—spanning 15 distinct agro-climatic zones—necessitates tailored microbial interventions aligned with regional cropping systems, soil chemistry, and farmer socioeconomic profiles (Table 2).

3.1 Revitalizing the Indo-Gangetic Plains and Curbing Stubble Burning

The Indo-Gangetic Plains (IGP), often termed India's breadbasket, suffer from severe soil fatigue induced by continuous rice–wheat monocropping and excessive chemical fertilizer application (Ladha et al., 2003; Bhattacharyya et al., 2015). Integrated Nutrient Management (INM) combining 75% recommended NPK with composite biofertilizers (*Azotobacter chroococcum*, phosphate-solubilizing *Bacillus megaterium*, *Rhizobium*, and potassium-solubilizing *Frateuria aurantia*) consistently sustains or increases grain yields while restoring soil microbial biomass carbon and dehydrogenase activity (Mader et al., 2011; Kumar et al., 2021; Kumar et al., 2024).

Agro-Climatic Region	**Dominant Cropping Systems**	**Key Soil & Climatic Constraints**	**Target Microbial Interventions**	**Documented Agronomic & Ecological Benefits**
Indo-Gangetic Plains (IGP)	Rice–wheat, rice–maize, sugarcane	Declining SOC, alkaline pH (7.8–8.5), P fixation, groundwater depletion, crop residue burning	*Azotobacter* + PSB + KSB consortia; *Trichoderma* & lignocellulolytic fungi (*Aspergillus*, *Phanerochaete*) for in situ straw decomposition	15–25% reduction in synthetic NPK; rapid 20–25 day paddy straw biodecomposition, curbing stubble burning
Rainfed & Semi-Arid Deccan / Central India	Pulses (chickpea, pigeonpea), oilseeds (groundnut, soybean), cotton, millets	Moisture deficit, terminal drought, low soil organic matter, *Fusarium* wilt, dry root rot	Native drought-tolerant *Rhizobium* + AMF (*F. mosseae*) + ACC-deaminase *Pseudomonas* + *T. harzianum*	12–28% yield gain under rainfed conditions; enhanced nodulation, P/Zn uptake, and wilt suppression
North-Eastern & Himalayan Hill Regions	Upland rice, maize, ginger, turmeric, citrus, tea, vegetables	Strong soil acidity (pH 4.5–5.5), Al/Fe toxicity, P immobilization, steep slope erosion	Acid-tolerant PSB (*Burkholderia*, *Penicillium*), AMF, *Azospirillum*, and indigenous *Trichoderma* strains	Mitigates Al toxicity; unlocks Fe/Al-bound P; supports certified organic farming (e.g., Sikkim model)
Coastal & Arid Salt-Affected Belts	Salt-tolerant rice, cotton, mustard, pearl millet, horticulture	Primary/secondary soil salinity and sodicity (ECe greater than 4 dS/m), high Na^+ toxicity	Halotolerant PGPR (*Bacillus licheniformis*, *Halomonas*, *Pseudomonas*) + salt-adapted AMF	Improves K^+/Na^+ selectivity, osmolyte accumulation, and grain yield by 15–30% on marginal saline lands

Beyond nutrient supply, microbial consortia offer an immediate solution to the acute environmental crisis of paddy straw burning in Punjab, Haryana, and western Uttar Pradesh. Application of lignocellulolytic microbial decomposers—such as the Pusa Decomposer consortium comprising *Aspergillus awamori*, *Trichoderma viride*, *Phanerochaete chrysosporium*, and *Aspergillus nidulans*—accelerates in situ breakdown of silica-rich rice residues within 20–25 days, recycling ~6–7 kg N, 1–1.5 kg P, and 15–20 kg K per tonne of straw back into the soil while eliminating particulate air pollution (Kumar et al., 2008; Chaudhary et al., 2017).

3.2 Climate-Proofing Rainfed Drylands and Pulse-Oilseed Systems

Over 80% of India's pulses, oilseeds, and coarse cereals (sorghum, pearl millet, finger millet) are cultivated in rainfed drylands characterized by erratic rainfall and impoverished soils (Venkateswarlu & Shanker, 2009). Dual inoculation of legumes such as chickpea (*Cicer arietinum*), pigeonpea (*Cajanus cajan*), and green gram (*Vigna radiata*) with host-specific *Mesorhizobium*/*Bradyrhizobium*/*Rhizobium*, phosphate-solubilizing bacteria, and arbuscular mycorrhizal fungi (*Funneliformis mosseae*) generates strong tripartite synergy: AMF improve phosphorus and water supply required for energy-intensive nitrogenase activity, while rhizobia and PSB enhance root colonization, phosphorus bioavailability, and nitrogen nutrition, boosting grain yields by 15–30% even under moisture-limited regimes (Mortimer et al., 2008; Teja et al., 2022; Calderon & Dangi, 2024; Mugabo et al., 2024). Furthermore, seed biopriming of rainfed crops with ACC-deaminase-producing *Pseudomonas* and *Trichoderma harzianum* enhances seedling vigor, deep rooting, and tolerance against dry root rot (*Macrophomina phaseolina*) (Glick et al., 2007; Harman et al., 2004).

3.3 Advancing Organic and Natural Farming in the Himalayas and Tribal Belts

North-Eastern India and Himalayan hill states possess inherently high soil organic matter and low baseline chemical fertilizer usage, positioning them as natural hubs for certified organic agriculture—exemplified by Sikkim's transition to 100% organic farming (Das et al., 2017). However, strong soil acidity (pH 4.5–5.5) in these regions causes severe phosphorus fixation by iron and aluminum oxides and aluminum phytotoxicity. Bioprospecting indigenous acid-tolerant phosphate-solubilizing bacteria (*Burkholderia*, *Pantoea*), acid-adapted AMF, and native *Trichoderma* isolates from Himalayan rhizospheres unlocks fixed phosphorus and protects high-value spice and horticultural crops (ginger, turmeric, large cardamom, mandarin) from devastating rhizome rot and wilt diseases without synthetic agrochemicals (Pandey et al., 2006; Negi et al., 2025).

Similarly, microbial science provides an empirical foundation to validate and optimize traditional Indian agroecological practices—such as *Jeevamrutha* and *Beejamrutha* promoted under Zero Budget Natural Farming (ZBNF) / Bharatiya Prakritik Krishi Paddhati (BPKP). While traditional fermented formulations introduce diverse native *Bacillus*, *Pseudomonas*, lactic acid bacteria, and yeasts that stimulate soil enzymatic activity, scientific standardization and enrichment with verified elite PGPR/BCA strains can transform these low-cost on-farm preparations into reliable, high-potency bio-inputs for resource-poor tribal and marginal farmers (Smith et al., 2020; Negi et al., 2025).

3.4 Biofortification and Phytoremediation of Degraded Lands

Micronutrient deficiencies—particularly zinc and iron—affect both crop productivity across ~36–49% of Indian soils and human nutrition in rural populations (Alloway, 2009). Zinc-solubilizing bacteria (*Bacillus aryabhattai*, *Pseudomonas* spp.) and AMF mobilize insoluble zinc compounds (ZnO, $ZnCO_3$) via organic acid secretion and enhance root-to-grain translocation, achieving agronomic biofortification of wheat, rice, and maize grains alongside higher yields (Ramesh et al., 2014; Singh & Prasanna, 2020). Additionally, heavy-metal-tolerant PGPR and mycorrhizae immobilize or detoxify arsenic, cadmium, and chromium in contaminated peri-urban and alluvial soils (such as arsenic-affected groundwater belts of West Bengal and Bihar) through biosorption, siderophore chelation, and altered redox speciation, safeguarding rural food chains (Saha et al., 2016; Glick, 2010).

4. Key Challenges Hindering Field-Scale Translation in Rural India

Despite overwhelming laboratory and greenhouse evidence, the transition of plant–microbe technologies from research stations to millions of smallholder fields in rural India faces multi-dimensional bottlenecks spanning biology, manufacturing, regulation, and extension (Herrmann & Lesueur, 2013; Bashan et al., 2014; Kumar et al., 2021).

4.1 Ecological Inconsistency and the "Lab-to-Field" Performance Gap

A major scientific barrier is the poor reproducibility of single-strain microbial inoculants across diverse field environments. Strains selected under sterile, nutrient-controlled laboratory conditions frequently fail to establish in complex native soils due to intense competition with indigenous microflora, predation by bacterivorous protozoa and bacteriophages, incompatible soil pH, high ambient temperatures, or mismatched host crop genotypes (Bulgarelli et al., 2013; Sessitsch et al., 2019). Moreover, excessive chemical fertilizer application in intensively farmed regions actively suppresses symbiotic signaling: high soil nitrate inhibits legume nodulation, while elevated soluble phosphate downregulates strigolactone exudation and reduces plant carbon allocation to AMF, rendering inoculants agronomically redundant when applied alongside full chemical doses (Parniske, 2008; Calderon & Dangi, 2024).

4.2 Formulation, Shelf Life, and Tropical Supply Chain Constraints

Unlike synthetic chemicals, biofertilizers and biopesticides contain living cells or propagules whose viability declines rapidly under rural Indian storage and transport conditions, where ambient temperatures routinely exceed 35–45 °C without cold-chain infrastructure (Bashan et al., 2014). Historically, Indian biofertilizer production relied heavily on solid lignite, charcoal, or talc carriers, which suffer from high contamination rates, batch-to-batch physicochemical variability, short shelf life (often less than 3–6 months), and difficulty in mechanized seed or drip application (Herrmann & Lesueur, 2013; Malusá et al., 2012). While liquid biofertilizers formulated with osmoprotectants (trehalose, glycerol, polyvinylpyrrolidone) have improved viability to 12–18 months, maintaining high colony-forming unit (CFU) counts ($10^8\text{–}10^9\text{ CFU/mL}$) during last-mile village retail distribution remains a persistent logistical hurdle (Kumar et al., 2021; Negi et al., 2025).

4.3 Quality Control Deficits and Regulatory Fragmentation

Farmer trust is the single most fragile asset in rural input markets. Substandard commercial products—containing low viable cell counts, dead inocula, or pathogenic contaminants—have periodically flooded local markets, causing field failures that disillusion farmers and trigger reversion to chemical inputs (Herrmann & Lesueur, 2013; Vaidyanathan, 2022). Although biofertilizers are regulated under India's Fertilizer (Inorganic, Organic or Mixed) Control Order (FCO), 1985, and biopesticides under the Insecticides Act, 1968 (Central Insecticides Board and Registration Committee, CIBRC), enforcement capacity at the district level is severely constrained. State soil and biofertilizer testing laboratories often lack molecular identification facilities (such as 16S rRNA/ITS sequencing or qPCR) to distinguish active functional strains from inert contaminants, relying instead on slow, culture-dependent plate counts (Vaidyanathan, 2022). Furthermore, multi-strain microbial consortia and dual-purpose products (exhibiting both biofertilizer and biopesticidal traits) encounter overlapping, protracted regulatory approval pathways that discourage private and startup innovation (Negi et al., 2025).

4.4 Socioeconomic, Behavioral, and Extension Barriers

Smallholder farmers in rural India exhibit strong risk aversion, as a single crop failure can precipitate severe household debt. Chemical fertilizers deliver rapid, visible vegetative greening within days, whereas microbial inoculants act gradually through root colonization and soil biological activation, yielding subtler early visual cues (Bashan et al., 2014; Kumar et al., 2021). Additionally, massive public subsidies on urea distort market price signals, making synthetic nitrogen artificially cheap compared to unsubsidized quality biostimulants (Sapkota & Bijay-Singh, 2025). Compounding this economic asymmetry is an acute last-mile knowledge gap: many farmers and local agri-input dealers lack training on proper seed inoculation techniques, compatibility with chemical seed treatments (e.g., fungicides that kill *Rhizobium* or *Trichoderma* upon direct contact), soil moisture requirements at sowing, and expiry verification (Herrmann & Lesueur, 2013).

5. Future Perspectives and Strategic Roadmap

Overcoming these constraints requires converging frontier microbiome science with frugal formulation engineering, decentralized rural biomanufacturing, and coherent policy reform.

5.1 Multi-Omics-Guided Synthetic Microbial Communities (SynComs)

Rather than relying on isolated single strains, the future of agricultural microbiome engineering lies in designing Synthetic Microbial Communities (SynComs)—rationally assembled, functionally complementary consortia combining nitrogen fixers, P/K solubilizers, AMF, and biocontrol agents that mimic natural core microbiomes (Vorholt et al., 2017; Sessitsch et al., 2019). Integrating high-throughput amplicon sequencing, shotgun metagenomics, metatranscriptomics, and metabolomics enables researchers to map indigenous rhizosphere networks across India's agro-climatic zones and identify keystone taxa adapted to local soil pH, moisture, and crop cultivars (Bulgarelli et al., 2013; Sahil et al., 2025). Machine learning and genome-scale

metabolic modeling can predict cross-feeding synergies and ecological stability prior to field deployment, ensuring that introduced SynComs coexist harmoniously with native soil microbiota and persist across the crop lifecycle (Vorholt et al., 2017; Gupta et al., 2021).

5.2 Host-Microbiome Co-Breeding ("Microbiome-Smart" Crops)

Modern crop breeding during the Green Revolution inadvertently selected for high yield under luxury synthetic fertilizer regimes, often eroding wild plant traits governing root exudation and beneficial microbial recruitment (Pérez-Jaramillo et al., 2016). Future Indian crop improvement programs—leveraging genome-wide association studies (GWAS) and marker-assisted selection—must incorporate "microbiome-recruitment traits" (such as optimal flavonoid and strigolactone exudation profiles and root hair architecture) into elite varieties of rice, wheat, pulses, and millets (Pérez-Jaramillo et al., 2016; Gupta et al., 2020). Breeding cultivars that actively nurture beneficial rhizosphere partners maximizes biological nutrient acquisition and stress resilience without requiring repeated high-dose inoculation.

5.3 Next-Generation Bioformulations: Biofilms, Nano-Encapsulation, and Seed Coating

To conquer tropical shelf-life and field survival barriers, formulation science must transition from crude carrier powders to advanced delivery matrices (Bashan et al., 2014; Malusá et al., 2012). Biofilm-based biofertilizers—wherein beneficial bacteria (e.g., *Azotobacter* or *Bacillus*) are co-cultured with fungal partners (*Trichoderma* or *Aspergillus*) to form robust, extracellular polymeric matrix-embedded communities—exhibit superior desiccation tolerance, UV resistance, and root colonization efficiency compared to planktonic cells (Seneviratne et al., 2008; Kumar et al., 2021). Similarly, **polymeric micro- and nano-encapsulation** using biodegradable biopolymers (sodium alginate, chitosan, starch, carboxymethyl cellulose) enriched with trehalose or humic acids shields microbes from high storage temperatures and agrochemical toxicity while enabling controlled, sustained release in the rhizosphere (Schoebitz et al., 2013; Kumar et al., 2021). Industrial and FPO-level **film-coating of seeds** with stable microbial consortia prior to sale eliminates on-farm application errors and guarantees uniform rhizosphere establishment from germination onward (O'Callaghan, 2016).

5.4 Decentralized Rural Biomanufacturing and Policy Alignment

Scaling microbial agriculture across 140 million Indian farm holdings cannot rely solely on centralized urban factories. Establishing decentralized, village- or block-level bio-input resource centers (BRCs) operated by Farmer Producer Organizations (FPOs), women's self-help groups (SHGs), and rural agri-entrepreneurs—supported by state agricultural universities (SAUs) and Krishi Vigyan Kendras (KVKs) providing certified mother cultures and quality testing—can dramatically reduce cold-chain costs, create rural green employment, and supply fresh, region-specific inoculants (Vaidyanathan, 2022; Sapkota & Bijay-Singh, 2025).

At the national policy level, integrating biofertilizer recommendations into India's Soil Health Card (SHC) scheme and digital public infrastructure (AgriStack) can shift farmers from blanket chemical applications to precision Integrated Nutrient

Management (Sapkota & Bijay-Singh, 2025). Reforming fertilizer subsidy architectures—such as introducing direct benefit transfers (DBT) or nutrient-based incentives that reward soil organic carbon accrual and verified biofertilizer adoption—will level the economic playing field. Simultaneously, mandating QR-code-based batch traceability, molecular quality certification, and harmonized single-window regulatory clearance for multi-functional microbial consortia will safeguard product efficacy and build enduring farmer confidence (Vaidyanathan, 2022; Negi et al., 2025).

6. Conclusion

Harnessing beneficial plant–microbe interactions represents a cornerstone strategy for reconciling agricultural productivity, environmental stewardship, and smallholder prosperity in rural India. Decades of mechanistic research confirm that PGPR, nitrogen-fixing diazotrophs, arbuscular mycorrhizal fungi, and biocontrol fungi can substantially reduce synthetic fertilizer and pesticide requirements while buffering crops against drought, salinity, and soil-borne diseases. Realizing this potential at scale, however, demands bridging the persistent gap between laboratory discovery and rural field reality. By uniting multi-omics-guided SynCom design and host-microbiome co-breeding with heat-stable polymeric and biofilm formulations, decentralized FPO-led biomanufacturing, rigorous molecular quality enforcement, and subsidy realignment, India can transition from chemically intensive farming to a resilient, regenerative, and self-reliant rural agricultural economy.

References

Abrol, Y. P., Adhya, T. K., Aneja, V. P., Raghuram, N., Pathak, H., Kulshrestha, U., Sharma, C., & Singh, B. (2017). *The Indian Nitrogen Assessment: Sources of Reactive Nitrogen, Environmental and Climate Effects, Management Options, and Policies*. Elsevier.

Alloway, B. J. (2009). Soil factors associated with zinc deficiency in crops and humans. *Environmental Geochemistry and Health*, 31(5), 537–548.

Alori, E. T., Glick, B. R., & Babalola, O. O. (2017). Microbial phosphorus solubilization and its potential for use in sustainable agriculture. *Frontiers in Microbiology*, 8, 971.

Arkhipova, T. N., Veselov, S. U., Melentiev, A. I., Martynenko, E. V., & Kudoyarova, G. R. (2005). Ability of bacterium *Bacillus subtilis* to produce cytokinins and to influence the growth and endogenous hormone content of lettuce plants. *Plant and Soil*, 272(1), 201–209.

Auge, R. M. (2001). Water relations, drought and vesicular-arbuscular mycorrhizal symbiosis. *Mycorrhiza*, 11(1), 3–42.

Bais, H. P., Weir, T. L., Perry, L. G., Gilroy, S., & Vivanco, J. M. (2006). The role of root exudates in rhizosphere interactions with plants and other organisms. *Annual Review of Plant Biology*, 57, 233–266.

Bano, A., & Fatima, M. (2009). Salt tolerance in *Zea mays* (L.) following inoculation with *Rhizobium* and *Pseudomonas*. *Biology and Fertility of Soils*, 45(4), 405–413.

Bashan, Y., de-Bashan, L. E., Prabhu, S. R., & Hernandez, J. P. (2014). Advances in plant growth-promoting bacterial inoculant technology: formulations and practical perspectives (1998–2013). *Plant and Soil*, 378(1–2), 1–33.

Bharti, N., Pandey, S. S., Barnawal, D., Patel, V. K., & Kalra, A. (2016). Plant growth promoting rhizobacteria *Dietzia natronolimnaea* modulates the expression of stress responsive genes providing protection of wheat from salinity stress. *Scientific Reports*, 6, 34768.

Bhattacharyya, P. N., & Jha, D. K. (2012). Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture. *World Journal of Microbiology and Biotechnology*, 28(4), 1327–1350.

Bhattacharyya, R., Ghosh, B. N., Mishra, P. K., Mandal, B., Rao, C. S., Sarkar, D., Das, K., Anil, K. S., Lalitha, M., Hati, K. M., & Franzluebbers, A. J. (2015). Soil degradation in India: Challenges and potential solutions. *Sustainability*, 7(4), 3528–3570.

Bijay-Singh, & Craswell, E. (2021). Fertilizers and nitrate pollution of surface and ground water: an increasingly pervasive global problem. *SN Applied Sciences*, 3(4), 518.

Birthal, P. S., Negi, D. S., Khan, M. T., & Agarwal, S. (2015). Is Indian agriculture becoming resilient to droughts? Evidence from rice production systems. *Food Policy*, 56, 1–12.

Boddey, R. M., Urquiaga, S., Alves, B. J., & Reis, V. (2003). Endophytic nitrogen fixation in sugarcane: present knowledge and future applications. *Plant and Soil*, 252(1), 139–149.

Bulgarelli, D., Schlaeppi, K., Spaepen, S., Ver Loren van Themaat, E., & Schulze-Lefert, P. (2013). Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant Biology*, 64, 807–838.

Calderon, R. B., & Dangi, S. R. (2024). Arbuscular mycorrhizal fungi and *Rhizobium* improve nutrient uptake and microbial diversity relative to dryland site-specific soil conditions. *Microorganisms*, 12(4), 667.

Chaudhary, S., Dheri, G. S., & Brar, B. S. (2017). Long-term effects of NPK fertilizers and organic manures on carbon stabilization and management index under rice-wheat cropping system. *Soil and Tillage Research*, 166, 59–66.

Compant, S., Duffy, B., Nowak, J., Clément, C., & Barka, E. A. (2005). Use of plant growth-promoting bacteria for biocontrol of plant diseases: principles, mechanisms of action, and future prospects. *Applied and Environmental Microbiology*, 71(9), 4951–4959.

Das, A., Patel, D. P., Kumar, M., Ramkrushna, G. I., Mukherjee, A., Layek, J., Ngachan, S. V., & Buragohain, J. (2017). Impact of seven years of organic farming on soil and produce quality and crop yields in eastern Himalayas, India. *Agriculture, Ecosystems & Environment*, 236, 142–153.

Dodd, I. C., & Pérez-Alfocea, F. (2012). Microbial amelioration of crop salinity stress. *Journal of Experimental Botany*, 63(9), 3415–3428.

Druzhinina, I. S., Seidl-Seiboth, V., Herrera-Estrella, A., Horwitz, B. A., Kenerley, C. M., Monte, E., Mukherjee, P. K., Zeilinger, S., Grigoriev, I. V., & Kubicek, C. P. (2011). *Trichoderma*: the genomics of opportunistic success. *Nature Reviews Microbiology*, 9(10), 749–759.

Glick, B. R. (2010). Using soil bacteria to facilitate phytoremediation. *Biotechnology Advances*, 28(3), 367–374.

Glick, B. R. (2012). Plant growth-promoting bacteria: mechanisms and applications. *Scientifica*, 2012, 963401.

Glick, B. R. (2014). Bacteria with ACC deaminase can promote plant growth and help to feed the world. *Microbiological Research*, 169(1), 30–39.

Glick, B. R., Todorovic, B., Czarny, J., Cheng, Z., Duan, J., & McConkey, B. (2007). Promotion of plant growth by bacterial ACC deaminase. *Critical Reviews in Plant Sciences*, 26(5–6), 227–242.

Gupta, R., Anand, G., Gaur, R., & Yadav, D. (2021). Plant–microbiome interactions for sustainable agriculture: a review. *Physiology and Molecular Biology of Plants*, 27(1), 165–179.

Gupta, S., Schillaci, M., Walker, R., Smith, P. M., Watt, M., & Roessner, U. (2020). Alleviation of salinity stress in plants by endophytic plant-fungal symbiosis: Current knowledge, perspectives and future directions. *Plant and Soil*, 461(1–2), 219–244.

Gururani, M. A., Upadhyaya, C. P., Baskar, V., Venkatesh, J., Nookaraju, A., & Park, S. W. (2013). Plant growth-promoting rhizobacteria enhance abiotic stress tolerance in *Solanum tuberosum* through inducing changes in the expression of ROS-scavenging enzymes and improved photosynthetic performance. *Journal of Plant Growth Regulation*, 32(2), 245–258.

Haas, D., & Défago, G. (2005). Biological control of soil-borne pathogens by fluorescent pseudomonads. *Nature Reviews Microbiology*, 3(4), 307–319.

Harman, G. E., Howell, C. R., Viterbo, A., Chet, I., & Lorito, M. (2004). *Trichoderma* species—opportunistic, avirulent plant symbionts. *Nature Reviews Microbiology*, 2(1), 43–56.

Herridge, D. F., Peoples, M. B., & Boddey, R. M. (2008). Global inputs of biological nitrogen fixation in agricultural systems. *Plant and Soil*, 311(1–2), 1–18.

Herrmann, L., & Lesueur, D. (2013). Challenges of formulation and quality of biofertilizers for successful inoculation. *Applied Microbiology and Biotechnology*, 97(20), 8859–8873.

Kaushal, M., & Wani, S. P. (2016). Plant-growth-promoting rhizobacteria: drought stress alleviators to ameliorate crop production in drylands. *Annals of Microbiology*, 66(1), 35–42.

Kloepper, J. W., Leong, J., Teintze, M., & Schroth, M. N. (1980). Enhanced plant growth by siderophores produced by plant growth-promoting rhizobacteria. *Nature*, 286(5776), 885–886.

Köhl, J., Kolnaar, R., & Ravensberg, W. J. (2019). Mode of action of microbial biological control agents against plant diseases: relevance beyond efficacy. *Frontiers in Plant Science*, 10, 845.

Kumar, R., Gaind, S., & Nain, L. (2008). Evaluation of thermophilic fungal consortium for paddy straw composting. *Biodegradation*, 19(3), 395–402.

Kumar, S., Diksha, Sindhu, S. S., & Kumar, R. (2021). Biofertilizers: An ecofriendly technology for nutrient recycling and environmental sustainability. *Current Research in Microbial Sciences*, 3, 100094.

Kumar, S. P., Singh, N., Kishore, A., et al. (2024). Effect of integrated nutrient management on growth and yield of green gram [*Vigna radiata* (L) Wilczek]. *Journal of Food Legumes*, 37(1), 114–116.

Ladha, J. K., Dawe, D., Pathak, H., Padre, A. T., Yadav, R. L., Singh, B., Singh, Y., Singh, Y., Singh, P., Kundu, A. L., Sakal, R., Ram, N., Regmi, A. P., Gami, S. K., Bhandari, A. L., Amin, R., Yadav, C. R., Bhattarai, E. M., Das, S., ... Hobbs, P. R. (2003). How extensive are yield declines in long-term rice–wheat experiments in Asia? *Field Crops Research*, 81(2–3), 159–180.

Lehmann, A., Veresoglou, S. D., Leifheit, E. F., & Rillig, M. C. (2014). Arbuscular mycorrhizal influence on zinc nutrition in crop plants—a meta-analysis. *Soil Biology and Biochemistry*, 69, 123–131.

Mader, P., Kaiser, F., Adholeya, A., Singh, R., Uppal, H. S., Sharma, A. K., Srivastava, R., Sahai, V., Aragno, M., Wiemken, A., Johri, B. N., & Fried, P. M. (2011). Inoculation of root microorganisms for sustainable wheat–rice and wheat–black gram rotations in India. *Soil Biology and Biochemistry*, 43(3), 609–619.

Malusá, E., Sas-Paszt, L., & Ciesielska, J. (2012). Technologies for beneficial microorganisms inocula used as biofertilizers. *The Scientific World Journal*, 2012, 491206.

Meena, V. S., Maurya, B. R., & Verma, J. P. (2014). Does a rhizospheric microorganism enhance K^+ availability in agricultural soils? *Microbiological Research*, 169(5–6), 337–347.

Mortimer, P. E., Pérez-Fernández, M. A., & Valentine, A. J. (2008). The role of arbuscular mycorrhizal colonization in the carbon and nutrient economy of the tripartite symbiosis with nodulated *Phaseolus vulgaris*. *Soil Biology and Biochemistry*, 40(5), 1019–1027.

Mugabo, J. P., Bhople, B. S., Kumar, A. K., Singh, V. K., Mahmoud, E. A., Ullah, F., Elansary, H. O., & Singh, O. (2024). Insights into the tripartite interaction: effects of Arbuscular mycorrhizae and *Rhizobium* on root morphology, soil enzymes, and biochemical properties in pea cultivation in alluvial soils of Punjab, India. *Cogent Food & Agriculture*, 10(1), 2366384.

Negi, R., Yadav, N., & Yadav, A. N. (2025). Microbial biofertilizers: A paradigm shift towards agricultural sustainability. *Biologia*, 80, 1–24.

O'Callaghan, M. (2016). Microbial inoculation of seed for improved crop performance: issues and opportunities. *Applied Microbiology and Biotechnology*, 100(13), 5729–5746.

Oldroyd, G. E. (2013). Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. *Nature Reviews Microbiology*, 11(4), 252–263.

Ongena, M., & Jacques, P. (2008). *Bacillus* lipopeptides: versatile weapons for plant disease biocontrol. *Trends in Microbiology*, 16(3), 115–125.

Pandey, A., Trivedi, P., Kumar, B., & Palni, L. M. S. (2006). Characterization of a phosphate solubilizing and antagonistic strain of *Pseudomonas putida* (B0) isolated from a sub-alpine location in the Indian Central Himalaya. *Current Microbiology*, 53(2), 102–107.

Parniske, M. (2008). Arbuscular mycorrhiza: the mother of plant root endosymbioses. *Nature Reviews Microbiology*, 6(10), 763–775.

Patten, C. L., & Glick, B. R. (2002). Role of *Pseudomonas putida* indoleacetic acid in development of the host plant root system. *Applied and Environmental Microbiology*, 68(8), 3795–3801.

Pérez-Jaramillo, J. E., Mendes, R., & Raaijmakers, J. M. (2016). Impact of plant domestication on rhizosphere microbiome assembly and functions. *Plant Molecular Biology*, 90(6), 635–644.

Pieterse, C. M., Zamioudis, C., Berendsen, R. L., Weller, D. M., Van Wees, S. C., & Bakker, P. A. (2014). Induced systemic resistance by beneficial microbes. *Annual Review of Phytopathology*, 52, 347–375.

Pingali, P. L. (2012). Green Revolution: Impacts, limits, and the path ahead. *Proceedings of the National Academy of Sciences*, 109(31), 12302–12308.

Ramesh, A., Sharma, S. K., Sharma, M. P., Yadav, N., & Joshi, O. P. (2014). Inoculation of zinc solubilizing *Bacillus aryabhattai* strains for improved growth, mobilization and biofortification of zinc in soybean and wheat cultivated in Vertisols of central India. *Applied Soil Ecology*, 73, 87–96.

Richardson, A. E., & Simpson, R. J. (2011). Soil microorganisms mediating phosphorus availability update on microbial phosphorus. *Plant Physiology*, 156(3), 989–996.

Rillig, M. C. (2004). Arbuscular mycorrhizae, glomalin, and soil aggregation. *Canadian Journal of Soil Science*, 84(4), 355–363.

Rodríguez, H., & Fraga, R. (1999). Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnology Advances*, 17(4–5), 319–339.

Saha, M., Sarkar, S., Sarkar, B., Sharma, B. K., Bhattacharjee, S., & Tribedi, P. (2016). Microbial siderophores and their potential applications: a review. *Environmental Science and*

Pollution Research, 23(5), 3984–3999.

Sahil, R., Pal, V., Kharat, A. S., & Jain, M. (2025). A multi-omics meta-analysis of rhizosphere microbiome reveals growth-promoting marker bacteria at different stages of legume development. *Plant, Cell & Environment*, 48(5), 3120–3138.

Sandhya, V., Ali, S. Z., Grover, M., Reddy, G., & Venkateswarlu, B. (2009). Alleviation of drought stress effects in sunflower seedlings by the exopolysaccharides producing *Pseudomonas putida* strain GAP-P45. *Biology and Fertility of Soils*, 46(1), 17–26.

Santi, C., Bogusz, D., & Franche, C. (2013). Biological nitrogen fixation in non-legume plants. *Annals of Botany*, 111(5), 743–767.

Sapkota, T. B., & Bijay-Singh. (2025). *Roadmap for Atmanirbharta in Fertilizers* (Policy Paper 150). National Academy of Agricultural Sciences (NAAS), New Delhi.

Sasse, J., Martinoia, E., & Northen, T. (2018). Feed your friends: do plant exudates shape the root microbiome? *Trends in Plant Science*, 23(1), 25–41.

Schoebitz, M., López, M. D., & Roldán, A. (2013). Bioencapsulation of microbial inoculants for better soil–plant fertilization: a review. *Agronomy for Sustainable Development*, 33(4), 751–765.

Seneviratne, G., Zavahir, J. S., Bandara, W. M. M. S., & Weerasekara, M. L. M. A. W. (2008). Fungal–bacterial biofilms: their development for novel biotechnological applications. *World Journal of Microbiology and Biotechnology*, 24(6), 739–743.

Sessitsch, A., Pfaffenbichler, N., & Mitter, B. (2019). Microbiome applications from lab to field: facing complexity. *Trends in Plant Science*, 24(3), 194–198.

Sharma, D. K., Singh, A., Sharma, P. C., Dagar, J. C., & Chaudhari, S. K. (2016). Sustainable management of sodic soils for crop production: opportunities and challenges. *Journal of Soil Salinity and Water Quality*, 8(2), 109–130.

Sharma, T. (2024). Antibiotic and antifungal characteristics of moringa (*Moringa oleifera*). *The Journal of Rural Advancement*, 12(1), 34–49.

Singh, D., & Prasanna, R. (2020). Potential of microbes in the biofortification of Zn and Fe in dietary food grains: A review. *Agronomy for Sustainable Development*, 40(2), 15.

Singh, J. S., Koushal, S., Kumar, A., Vimal, S. R., & Gupta, V. K. (2016). Book review: Microbial inoculants in sustainable agricultural productivity—Vol. II: Functional application. *Frontiers in Microbiology*, 7, 2105.

Smith, J., Yeluripati, J., Smith, P., & Nayak, D. R. (2020). Potential yield challenges to scale-up of zero budget natural farming. *Nature Sustainability*, 3(3), 247–252.

Smith, S. E., & Read, D. J. (2008). *Mycorrhizal Symbiosis* (3rd ed.). Academic Press.

Spaepen, S., Vanderleyden, J., & Remans, R. (2007). Indole-3-acetic acid in microbial and microorganism-plant signaling. *FEMS Microbiology Reviews*, 31(4), 425–448.

Teja, M. S., Kishore, A., Sharma, J. D., Kumar, S. P., Reddy, K. S., & Pramod, T. (2022). Different levels of phosphorus and phosphorus solubilizing bacteria (PSB) influence growth, yield and economics of green gram [*Vigna radiata* L.]. *Journal of Food Legumes*, 35(3), 180–183.

Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R., & Polasky, S. (2002). Agricultural sustainability and intensive production practices. *Nature*, 418(6898), 671–677.

Tyśkiewicz, R., Nowak, A., Ozimek, E., & Jaroszuk-Ściseł, J. (2022). *Trichoderma*: The current status of its application in agriculture for the biocontrol of fungal phytopathogens and stimulation of plant growth. *International Journal of Molecular Sciences*, 23(4), 2329.

Vaidyanathan, S. (2022). *State of Biofertilizers and Organic Fertilizers in India*. Centre for Science and Environment (CSE), New Delhi.

Van Wees, S. C., Van der Ent, S., & Pieterse, C. M. (2008). Plant immune responses triggered by beneficial microbes. *Current Opinion in Plant Biology*, 11(4), 443–448.

Vandenkoornhuyse, P., Quaiser, A., Duhamel, M., Le Van, A., & Dufresne, A. (2015). The importance of the microbiome of the plant holobiont. *New Phytologist*, 206(4), 1196–1206.

Venkateswarlu, B., & Shanker, A. K. (2009). Climate change and agriculture: adaptation and mitigation strategies. *Indian Journal of Agronomy*, 54(2), 226–230.

Vessey, J. K. (2003). Plant growth promoting rhizobacteria as biofertilizers. *Plant and Soil*, 255(2), 571–586.

Vorholt, J. A., Vogel, C., Carlström, C. I., & Müller, D. B. (2017). Establishing causality: opportunities of synthetic communities for plant microbiome research. *Cell Host & Microbe*, 22(2), 142–155.

Yang, J., Kloepper, J. W., & Ryu, C. M. (2009). Rhizosphere bacteria help plants tolerate abiotic stress. *Trends in Plant Science*, 14(1), 1–4.

Zhang, H., Kim, M. S., Sun, Y., Dowd, S. E., Shi, H., & Paré, P. W. (2008). Soil bacteria confer plant salt tolerance by tissue-specific regulation of the sodium transporter *HKT1*. *Molecular Plant-Microbe Interactions*, 21(6), 737–744.