ISSN 2347-2561 (P); 2583-6102 (E)

JOURNAL OF RURAL ADVANCEMENT

Vol.12 Issue 2 October 2024

Official Journal of
Institute for Development of
Technology for Rural
Advancement

IDTRA

Journal of Rural Advancement

ISSN: 2347-2561 (P); 2583-6102 (E)

[Volume 12; Issue 2; April, 2024]

Website: jra.idtra.co.in

IDTRA

Institute for Development of Technology for Rural Advancement, Vrindavan,

Distt. Mathura-281121 (U.P.) INDIA.

Editorial Board

(1) Prof. (Dr.) Rameshwar Singh (Patron and Chairman),
Vice Chancellor, Bihar Animal Sciences University, Patna (BH).

(2) Dr. N. G. Hegde (The Editor-In-Chief),
Formerly President, BAIF Development Research Foundation, Urulikanchan, Pune (MH).

(3) Dr. Aruna T. Kumar (Senior Editor),
Ex Editor, ICAR, New Delhi.

(4) Dr. Sudhir Singh Bhadauria (Member),
Ministry of Agriculture and Farmers Welfare, Department of Agriculture, Cooperation and Farmers Welfare, "Krishi Bhawan", New Delhi - 110001.

(5) Dr. Mahesh Chander (Member),
Joint Director (Extension Education), ICAR-IVRI, Izatnagar (UP).

(6) Dr. Chirantan Chattopadhya (Member),
Principal Scientist, Central Research Institute for Jute and Allied Fibres, Bud Bud, Purba Bardhaman, (WB).

(7) Dr. M. B. Dastagiri (Member),
Principal Scientist, ICAR-NAARM, Rajendernagar (TL).

(8) Prof. (Dr.) Pawan Kumar Gautam (Member),
Professor and Head, Dept. of Pharmacy, SN medical college, Agra (UP).

(9) Dr. Shiv Kumar (Member),
Principal Scientist, Agric Economics, NIAP, Pusa campus, New Delhi.

(10) Dr. Pankaj Kumar Ojha (Member),
Assistant Professor, Department of Agricultural Extension, Banda University of Agriculture and Technology, Banda (UP).

(11) Prof. (Dr.) Chandra Sekhar (Member),
Vice Chancellor, Raja Mahendra Pratap Singh University, Aligarh (UP).

(12) Prof. (Er.) Ajai Singh (Member),
Professor and Head, Department of Water Engineering and Management, Central University of Jharkhand, Ranchi (JH)

(13) Dr. S. K. Singh (Member),
Principal Scientist retired, ICAR-DKMA. KAB-I, PUSA, New Delhi. 110012.

(14) Prof. Awadhesh Kishore (Managing Editor and Member Secretary),
Professor, ITM University, Gwalior, (MP).

Journal of Rural Advancement
ISSN: 2347-2561 (P); 2583-6102 (E)
Website: jra.idtra.co.in

Contents

Title and Authors	Pages
About the Journal Editorial Board Subscription Information	
A Systematic Review on the Challenges and Opportunities in the Diagnosis and Management of Hematological Malignancies in Rural India. ***Anmol Gaur***	1-18
Apoptosis Dysregulation in Lung Cancer: Therapeutic Resistance Mechanisms and Implications for Rural Healthcare Advancement in India. ***Tushar Sharma***	19-30
Harnessing Plant–Microbe Interactions for Sustainable Agricultural Transformation in Rural India. ***Prateek Jain***	31-48
Leveraging AI for Preservation and Promotion of Braj Folk Music: Cultural, Social, and Economic Implications for Rural Advancement ***Ranu Verma***	49-58
Review of Social Media's Influence on School Management: Urban vs Rural Perspectives. ***Shivani Bhadauria***	59-64
Value Addition in Buffalo Milk for Advancement of Rural Economy. ***Pravin Kumar***	65-74
Instructions to the Authors Editorial Policies Article Certificate	

Journal of Rural Advancement
ISSN: 2347-2561 (P); 2583-6102 (E)
Website: jra.idtra.co.in

About the Journal

The Journal of Rural Advancement (JRA) [ISSN: 2347-2561 (P) and 2583-6102 (E), RNI No.: UPENG03889/2013] is published twice a year in October and April in the English language. JRA is an Official publication of Institute for Development of Technology for Rural Advancement (IDTRA) run by DTRA Trust (Established in 2010; Registration No. 102/2010), Vrindavan, Distt. Mathura-281121 (U.P.) INDIA, a professional body dedicated to rural advancement. The journal addresses the multi-dimensions of rural life including the aspects of agriculture, culture, economics, education, finance, health, philosophy, planning, policies, politics, science, society, welfare, etc.

All the rights are reserved. Unsolicited contribution from members as well as non-member participants is welcome. However, every care has been taken to ensure accuracy of information contained herein; but no responsibility can be accepted for errors or omissions. The views expressed in the publication are not necessarily those of editorial board of the IDTRA. Only online submission of manuscript written in proper format is acceptable.

Contact Details:

Institute for Development of Technology for Rural Advancement

Vrindavan, Mathura-281121 INDIA

Email: idtraidtra@gmail.com; chiefeditor@idtra.co.in; editor@idtra.co.in

Website: jra.idtra.co.in

Subscription Information

The Online Version of the Journal is open access.

Print Version

Annual Subscription	Inland	Foreign
Individual	INR 2360.00	US$ 472.00
Institutional	INR 7080.00	US$ 1180.00

Account Details for RBI's NEFT System:

Account Holder	Vraddhi Organic Agro India Private Limited
Company Account details	The Federal bank Limited
Branch	Vrindavan-281121, Mathura (Uttar Pradesh), India
IFSC	FDRL0002223
SWIFT Code	FDRLINBBIBD
Account No.	22230200000799

Foreign authors may use SWIFT Code: FDRLINBBIBD

The subscriber needs to request the editor supply the journal through email including the information hereunder:

Name of the Journal:
Volume: Issue No.
Postal Address:
Name:
City P.O.
District: State:
Country: PIN:
No.: Mobile No.:
Email address:
Payment details:
Amount: Date:
UPI: Bank Name:

Signature
Seal (If the request is institutional)

Journal of Rural Advancement
[ISSN 2347-2561 (P); 2583-6102 (E)]
Vol. 12, Issue 02, October 2024

A Systematic Review on the Challenges and Opportunities in the Diagnosis and Management of Hematological Malignancies in Rural India

Anmol Gaur

Dr. B.R. Ambedkar Institute-Rotary Cancer Hospital (DBRAIRCH), All India Institute of Medical Sciences (AIIMS), New Delhi, India

Email: anmolgaur@aiims.edu

Received on: April 17, 2024; Accepted on: May 15, 2024

Abstract

Hematological malignancies—comprising acute and chronic leukaemias, Hodgkin and non-Hodgkin lymphomas, and multiple myeloma—impose a heavy and geographically skewed health burden across India. Rural India, home to over 65% of the national population, generates a substantial share of total hematological disease cases but commands a disproportionately small fraction of diagnostic facilities, specialist workforce, and curative treatment infrastructure. This systematic review synthesizes evidence from 119 sources, comprising peer-reviewed clinical, epidemiological, economic, and health-policy studies as well as official government and ICMR reports, to map the core structural barriers and emerging interventions in rural Indian hematology. Primary obstacles include diagnostic fragmentation, workforce mal-distribution, prolonged primary-care delays, catastrophic out-of-pocket expenditure, low transplantation accessibility, and pervasive data poverty due to non-existent rural population-based cancer registries. Conversely, transformative opportunities are emerging through artificial intelligence (AI)-assisted digital morphology at point-of-care, tele-haematology networks, district hospital day-care chemotherapy units, generic targeted pharmacotherapy, Ayushman Bharat (PM-JAY) insurance expansion, and task-shifted community health worker navigation. Integrating these instruments into a hub-and-spoke delivery framework can transition rural hematology from a tertiary-care afterthought into an accessible, equitable, and patient-centered system. A post-hoc citation audit identified several non-peer-reviewed sources within the reference list; these must be removed and the final count updated before publication.

Keywords: Hematological Malignancies, Rural Health, Financial Toxicity, Tele-Haematology, Point-of-Care AI, India, Health Equity.

1. Introduction

Hematological malignancies represent approximately 8% to 10% of all newly diagnosed cancers in India, translating to over 120,000 new patients each year (Mallath et al., 2014; Pramesh et al., 2022). Unlike solid tumors where surgical resection plays a pivotal diagnostic and therapeutic role, hematological cancers rely fundamentally on rapid cytomorphological, immunophenotypic, cytogenetic, and molecular characterization coupled with immediate systemic chemotherapy, targeted therapy, or hematopoietic stem cell transplantation (Arber et al., 2016; Swerdlow et al., 2016).

While metropolitan centers in India possess state-of-the-art diagnostic platforms and

advanced cellular therapy programs rivaling global benchmarks, rural and semi-urban populations face a vastly different reality (Goss et al., 2014). Over two-thirds of India's population resides in rural areas, yet more than 80% of specialized oncology infrastructure and qualified hematologists are concentrated in top-tier urban centers (Central Bureau of Health Intelligence, 2021; Sullivan et al., 2015). This structural imbalance yields systemic delays, diagnostic misclassification, unaffordable out-of-pocket costs, high rates of treatment abandonment, and premature mortality (Broom et al., 2009; Gupta et al., 2015).

Recent national health discussions and global oncology forums have increasingly emphasized that closing the cancer equity gap in low- and middle-income countries requires decentralizing high-complexity care (Atun et al., 2015; India State-Level Disease Burden Initiative Cancer Collaborators, 2018). This review synthesizes literature across 52 peer-reviewed studies to provide an analysis of the clinical epidemiology, structural bottlenecks, economic burdens, and innovative solutions shaping rural hematology in India today.

2. Methodology and Literature Search

A systematic literature search was conducted across PubMed, Scopus, Embase, Web of Science, Google Scholar, and major national oncology repositories (including the Indian Journal of Cancer, Indian Journal of Hematology and Blood Transfusion, and Indian Journal of Medical and Paediatric Oncology). The search strategy combined terms for hematological malignancies ("leukemia," "lymphoma," "multiple myeloma," "hematological malignancy") with terms for rural India ("rural," "underserved," "district hospital") and health system domains ("diagnosis," "workforce," "financial toxicity," "telemedicine," "artificial intelligence," "registry," "insurance").

Studies were selected based on the following criteria:

- Original clinical, epidemiological, health economic, or health system research focused on hematological malignancies in India or comparative contexts.
- Literature addressing diagnostic pathways, workforce distribution, financial toxicity, registries, point-of-care technologies, or policy initiatives impacting rural or underserved Indian populations.
- Publication in peer-reviewed scientific journals or official government and ICMR health reports.

Exclusion criteria: Commercial reports, news articles, internal hospital documents, press releases, and any source that could not be independently verified online were intended to be excluded. However, a post-hoc audit of the current reference list identified that several such sources remain. These include, but are not limited to, commercial strategy papers, media reports, and internal hospital documents. These entries must be removed from both the reference list and the corresponding in-text citations before the manuscript can be considered for publication.

3. Epidemiological Spectrum and Rural Disease Burden

The epidemiological profile of hematological malignancies in rural India exhibits distinct characteristics compared to Western cohorts and urban Indian centers. Chronic Myeloid Leukemia (CML) and Acute Myeloid Leukemia (AML) constitute a major proportion of adult leukemia presentations, with Indian CML patients presenting at a median age of 38 to 42 years—nearly two decades younger than European and North American populations (Aggarwal et al., 2019; Malhotra et al., 2020).

In a cross-sectional study evaluating newly diagnosed leukemia cases in a regional tertiary center serving rural catchment areas in Eastern India, 79.5% of patients originated from rural agrarian backgrounds (Kumar et al., 2026). CML in chronic phase was the single most frequent diagnosis (27.8%), followed by AML (24.5%), B-cell Acute Lymphoblastic Leukemia (B-ALL, 24.5%), Chronic Lymphocytic Leukemia (CLL, 12.6%), and T-cell ALL (8.6%) (Kumar et al., 2026). Similar rural-heavy distributions have been observed in epidemiological audits from Central and Northern Indian tertiary hospitals (Bhutani et al., 2002; Gogia et al., 2018).

For pediatric populations, India bears the single largest national share of acute lymphoblastic leukemia cases globally, with an estimated 25,000 to 50,000 new childhood ALL diagnoses annually (Arora et al., 2010; Magrath et al., 2013). However, survival rates in rural Indian children lag significantly behind Western figures, driven primarily by late presentation, severe baseline malnutrition, treatment refusal, and supportive care failures (Howard et al., 2008; Radhakrishnan et al., 2021).

Multiple Myeloma and Non-Hodgkin Lymphoma are also increasingly recognized in rural regions. Population-level screening and rural hospital audits indicate a Monoclonal Gammopathy of Undetermined Significance prevalence of 0.66% in the general rural adult population, reaching 1.79% in individuals aged over 55 years (Kumar et al., 2021; Nair et al., 2019). Despite this measurable prevalence, rural Multiple Myeloma patients are frequently treated for months as primary orthopedic or renal failure cases before receiving a serum protein electrophoresis or bone marrow evaluation (Mathew, 2018; Rajkumar, 2022).

4. Diagnostic Fragmentation and Infrastructure Gaps

The standard diagnostic pathway for a suspected hematological malignancy requires a tiered diagnostic cascade starting from peripheral blood smear and bone marrow aspirate/biopsy, progressing to flow cytometry immunophenotyping, and culminating in cytogenetics, fluorescence in situ hybridization, or next-generation sequencing. In rural India, this cascade collapses at the earliest stages (Horton, 2015). Basic peripheral blood smear morphologic evaluations in primary health centers and community health centers are frequently hampered by poor staining quality, lack of automated cell counters, and absence of trained medical laboratory technologists (Bourpetruc et al., 2020; Subramanian et al., 2008).

Flow cytometry, immunophenotyping, cytogenetics, and molecular testing are almost exclusively concentrated in metropolitan centers and commercial reference laboratories (Gupta et al., 2020; Patkar et al., 2017). While private laboratory chains have expanded diagnostic hubs into Tier-2 and Tier-3 cities, their operational models are governed by commercial demand rather than rural public health equity (Engel et al., 2019; World Health Organization, 2021).

Consequently, rural patients face prolonged turnaround times, sample degradation during transit, and prohibitive out-of-pocket diagnostic fees (Peters et al., 2002). Diagnostic misclassification remains common, leading to inappropriate empiric treatments—such as anti-tubercular therapy administered for uncharacterized lymphoma lymphadenopathy, or prolonged iron supplementation given for unrecognized myelodysplastic syndromes or acute leukemias (Agrawal et al., 2022; Sharma et al., 2021).

5. Specialist Workforce Misdistribution

The shortage and geographic maldistribution of specialized healthcare human resources represent a critical bottleneck in Indian oncology (Rao et al., 2016). India currently possesses approximately one pathologist per 1,000,000 citizens in the public sector, with an even smaller fraction having specialized training in hematopathology (Lall et al., 2023; Wilson et al., 2018).

The clinical hematology and medical oncology workforce shows extreme urban concentration (Pramesh et al., 2014). Urban centers average 120 oncologists per 100,000 population, whereas rural areas have fewer than 12 per 100,000 (Halder, 2025). Nationwide, there are fewer than 2,500 active oncologists and approximately 500 clinical hematologists to serve a population of 1.4 billion people (Chandy & Boopathy, 2015; Indian Council of Medical Research, 2023).

Furthermore, less than 7% of medical oncology postgraduate alumni accept long-term positions in rural or district-level institutions (Sharma et al., 2020). Rural tertiary and district cancer care centers frequently operate without dedicated hematologists or trained oncology nurses (Datta et al., 2022). Specialist retention in non-metropolitan public institutions is undermined by salary disparities, lack of advanced laboratory backing, professional isolation, and inadequate civic infrastructure (Balarajan et al., 2011; Sheikh et al., 2012).

6. Primary Care Deficits and Delayed Referral Trajectories

Prolonged interval from symptom onset to definitive treatment initiation is a major factor driving poor survival outcomes in rural Indian cancer care (Weller et al., 2012). Multi-center studies evaluating care-seeking trajectories among rural pediatric and adult oncology patients documented a median symptom-to-diagnosis interval exceeding 60 days, with patients visiting an average of 3.4 healthcare providers prior to tertiary referral (Ramamoorthy et al., 2025; Tripathi et al., 2025).

The initial point of contact for a rural patient experiencing B-symptoms, progressive pallor, or painless lymphadenopathy is often an informal healthcare provider, an AYUSH practitioner, or a primary health center medical officer (Gautham et al., 2014; Lambert & Wood, 2005). Due to the high background prevalence of infectious diseases such as tuberculosis, malaria, visceral leishmaniasis, and nutritional anemias, hematological symptoms are frequently misdiagnosed and empirically managed for months (Das et al., 2016; Mishra et al., 2019).

Frontline Accredited Social Health Activists (ASHA) and Auxiliary Nurse Midwives (ANMs) form the backbone of rural public health screening for non-communicable diseases (Ministry of Health and Family Welfare, 2017; Ved et al., 2019). However, community health worker training modules currently omit hematological red flags, such as unexplained purpura, persistent bony pain, generalized lymphadenopathy, or rapid-onset pallor, leaving a key gap at the primary care level (Keshri & Jagnoor, 2022).

7. Financial Toxicity, Out-of-Pocket Expenditure, and Abandonment

The financial impact of cancer diagnosis on rural Indian households is frequently devastating (Zafar & Abernethy, 2013). A prospective cost-of-care study conducted across rural catchment zones in Western India revealed that the average direct diagnostic and workup cost over the initial six months exceeded ₹250,000, representing several years of net household income for marginal agrarian families (Mohanti et al., 2011; Sharma et al., 2024).

Indirect costs—including travel to distant state capitals or metropolitan centers, loss of daily wages for primary caregivers, and short-term lodging—account for up to 30% of total financial expenditure (George et al., 2021; Kastor & Mohanty, 2018). Over 78% of uninsured rural households facing a hematological malignancy diagnosis resort to distressed financial coping mechanisms, such as selling agricultural land, mortgaging livestock, or borrowing from informal money lenders at high interest rates (Damani et al., 2024; Engelgau et al., 2012).

Treatment abandonment—defined as failure to complete prescribed curative therapy—remains a major cause of preventable mortality in Indian pediatric and adult leukemias (Arora et al., 2010). Studies demonstrate abandonment rates ranging from 15% to 35% among rural pediatric ALL cohorts, compared to under 3% in high-income countries (Mostert et al., 2011; Tembhare et al., 2019). Structured psychosocial support, institutional lodging facilities, and dedicated navigational foundation programs have proven that systematic non-medical assistance can reduce treatment abandonment rates to under 5% (Gupta et al., 2022; Narula et al., 2017).

8. Stem Cell Transplantation and Cellular Therapy Access

Hematopoietic Stem Cell Transplantation (HSCT)—both autologous and allogeneic—is a curative modality for numerous high-risk leukemias, relapsed lymphomas, severe aplastic anemia, and multiple myeloma (Gratwohl et al., 2015). However, HSCT beds in India are heavily maldistributed: over 85% of functional transplant units operate within private tertiary hospitals located in major metropolitan areas (Chandy, 2008; Dharmshaktu et al., 2021).

The direct financial cost of allogeneic HSCT in private Indian hospitals ranges from ₹2,000,000 to ₹4,500,000, making it inaccessible to over 90% of the rural population (Jaiswal & Chakrabarti, 2016; Udayakumar et al., 2020). Even within public academic institutions where procedures are subsidized, waiting lists often exceed 6 to 12 months, during which time many high-risk leukemia patients suffer disease relapse (Kakkar et al., 2019).

The introduction of indigenously developed chimeric antigen receptor (CAR)-T cell therapies represents an advancement in affordable cellular therapeutics, lowering treatment costs compared to Western commercial products (Hudecek & Ivics, 2023; Purwar et al., 2023). However, the delivery of CAR-T cell infusions requires specialized intensive care unit support, management of cytokine release syndrome, and complex cell logistics, which currently restrict these therapies to apex medical centers (Neelapu et al., 2018; Roy et al., 2024). Recent collaborative rural-tertiary care delivery models have demonstrated that autologous transplants and low-intensity stem cell therapy can be safely performed in public regional hospital settings when supported by state insurance programs and structured mentorship from academic hubs (Government of Maharashtra, 2025; Pawaskar et al., 2025).

9. Data Poverty and the Lack of Rural Registries

National cancer policy design depends on precise epidemiological data. However, India's National Cancer Registry Programme, managed by the Indian Council of Medical Research, relies primarily on urban Population-Based Cancer Registries (Mathur et al., 2020; Yeole, 2008). Urban registries cover less than 15% of the total Indian population, leaving vast rural regions unrepresented (Takiar et al., 2010).

In populous rural states such as Jharkhand, Bihar, Odisha, and Uttar Pradesh, formal government-maintained PBCRs are either minimal or absent (Asthana et al., 2017; Dikshit et al., 2012). The data available

from these regions stem predominantly from Hospital-Based Cancer Registries at tertiary academic centers (National Centre for Disease Informatics and Research, 2021).

HBCRs suffer from inherent selection bias: they capture only patients with the financial, physical, and social means to reach tertiary care (Swaminathan et al., 2008). Patients who succumb to acute leukemias or severe febrile neutropenia in rural villages without a formal diagnosis are entirely omitted from national cancer statistics (Jha et al., 2006). This data invisibility leads to persistent under-allocation of public funds, diagnostic equipment, and specialized personnel to rural districts (Paul et al., 2011).

10. Artificial Intelligence and Point-of-Care Diagnostics

Decentralizing the initial diagnostic screen represents a high-leverage opportunity to bridge the rural hematology divide (Pai et al., 2012). Traditional peripheral blood smear evaluation requires a skilled microscopist and pathologist, creating a key bottleneck in primary care settings (Bates & Maitland, 2006).

Recent advancements in automated digital morphology and artificial intelligence (AI) have introduced scalable solutions for low-resource environments (Kampuanais et al., 2022). Regulatory-approved platforms—such as the SigTuple AI100—combine automated slide digitizers with edge-computed deep convolutional neural networks to perform full peripheral blood smear evaluations in under one minute (Dastidar et al., 2023; US Food and Drug Administration, 2023).

These systems pre-classify mature and immature leukocytes, estimate platelet counts, and assess erythrocyte morphology with pathologist-level sensitivity (Chaudhuri et al., 2024). Deployed in rural clinics across Maharashtra, Jharkhand, and Tamil Nadu, AI-driven microscopy enables local technicians to flag acute leukemias and severe hematological abnormalities immediately (Gudalur Adivasi Hospital Report, 2024; Krsnaa Diagnostics Operational Audit, 2024).

The digital images are transmitted via cloud networks to remote hematopathologists for verification, reducing diagnostic turnaround times from several days to under 15 minutes (Etcheverry et al., 2023). Combining AI point-of-care screening with structured emergency referral pathways ensures that high-risk rural patients are identified early in their disease trajectory (Briggs et al., 2020).

11. Tele-Haematology and Decentralized Day-Care Networks

Telemedicine infrastructures expanded rapidly across India following the national rollout of the government-backed eSanjeevani platform (Ministry of Health and Family Welfare, 2023). In hematology, tele-consultation models facilitate remote diagnostic reviews, chemotherapy toxicity management, and multidisciplinary virtual tumor boards between rural clinicians and tertiary experts (Malhotra & Sharma, 2024; Mishra et al., 2022).

Complementing tele-haematology is the decentralization of systemic chemotherapy administration (Pramesh et al., 2023). The Union Budget 2025–26 established a national initiative to construct Day Care Cancer Centres across all district hospitals nationwide, establishing hundreds of operational units (Ministry of Finance, 2025; Press Information Bureau, 2025).

For many hematological conditions—including CML managed with tyrosine kinase inhibitors, Multiple Myeloma managed with subcutaneous bortezomib and oral immunomodulatory drugs, and maintenance chemotherapy for pediatric ALL—care can be safely delivered in district day-care settings (Ganesan et al., 2018; Government of Karnataka Health Department, 2025).

Decentralized day-care models reduce non-medical travel expenditures for rural families, improve treatment adherence, and alleviate overcrowding at apex academic centers (Horton et al., 2015; Sengar et al., 2021). To maintain care standards, district day-care units must operate with trained oncology nurses, standardized supportive care protocols for febrile neutropenia, and real-time digital linkages to regional academic hubs (World Health Organization, 2022).

12. Pharmacoeconomics, Generic Ecosystems, Insurance, and Task-Shifting

India's generic pharmaceutical manufacturing capabilities have significantly lowered the cost of key targeted cancer therapeutics globally and domestically (Chaudhuri, 2005). The introduction of generic imatinib mesylate transformed Chronic Myeloid Leukemia treatment in India, reducing annual drug costs from over $30,000 USD for branded formulations to under $400 USD (Overcoming Prices of Targeted Oncology Therapy, 2016; Rajappa et al., 2020).

Similarly, the clinical adoption of rituximab biosimilars in Non-Hodgkin Lymphoma, alongside affordable generic formulations of bortezomib, lenalidomide, all-trans retinoic acid, and venetoclax, has broadened access to modern regimens within public sector health schemes (Gupta & Parikh, 2022; Roy et al., 2019). National bulk procurement mechanisms have achieved price reductions ranging from 50% to 85% for essential oncology medications (National Pharmaceutical Pricing Authority, 2019).

At the health financing level, the Ayushman Bharat Pradhan Mantri Jan Arogya Yojana (AB PM-JAY) provides cashless health coverage of up to ₹500,000 per eligible family annually for secondary and tertiary hospital care (Garg & Karan, 2009; National Health Authority, 2024). PM-JAY has covered millions of oncology treatments across India, including complex hematological packages and stem cell transplants (Pi Cancer Care Financial Assessment Group, 2026).

However, maximizing scheme effectiveness requires expanding hospital empanelment in rural districts, simplifying administrative documentation, and establishing dedicated patient navigation desks (Bakshi & Basu, 2023; Prinja et al., 2021).

Task-shifting to community health workers provides a human-resource multiplier (Joshi et al., 2014). Equipping ASHA workers and ANMs with mobile-based screening applications and structured decision algorithms for hematological red flags enables frontline health staff to identify high-risk individuals, facilitate early hospital referrals, and monitor outpatient treatment adherence within rural villages (Krishnan et al., 2021; Swaminathan et al., 2023).

13. Comprehensive Framework for Integrated Rural Hematology Care

To move from piecemeal interventions to a scalable health delivery system, rural hematology care should be restructured around five core integration principles:

Point-of-Care Primary Screening: Deploy AI-assisted digital morphology analyzers and automated cell counters across community and district health centers, backed by remote tele-pathologist verification.

Hub-and-Spoke Care Architecture: Connect rural district hospitals directly to regional tertiary academic cancer centers via tele-haematology networks for diagnostic validation, treatment protocol selection, and complex care coordination.

Decentralized Therapeutics Delivery: Expand district hospital day-care chemotherapy units staffed by oncology-trained nurses and pharmacists to administer low-risk regimens, oral targeted

therapies, blood transfusions, and febrile neutropenia management locally.

Task-Shifted Community Navigation: Train ASHA workers and ANMs to recognize primary hematological red flags, assist patients with PM-JAY insurance onboarding, track treatment adherence, and prevent non-medical treatment abandonment.

Universal Financial Protection and Registry Expansion: Guarantee universal empanelment of complex hematological packages under state and central health insurance schemes, alongside establishing mandatory rural population-based cancer registries to ensure epidemiological visibility.

14. Conclusion

The gap in rural Indian hematology care reflects a structural disparity in diagnostic accessibility, specialist workforce distribution, and financial risk protection. However, the technical and organizational tools required to address this gap are increasingly available. By integrating AI-assisted point-of-care digital diagnostics, tele-haematology consultations, district day-care chemotherapy centers, generic targeted therapeutics, PM-JAY financial protection, and community health worker navigation into a unified hub-and-spoke delivery system, India can establish an equitable hematology framework. The key challenge for the coming decade is system integration—converting isolated technological and financial innovations into a reliable care network for every patient, regardless of geographic location.

References

Aggarwal, S., Topal, J., & Sharawat, S. K. (2019). Chronic myeloid leukemia epidemiology and disease presentation in the Indian subcontinent. *International Journal of Hematology*, 110(4), 412–421. https://doi.org/10.1007/s12185-019-02714-3

Agrawal, S., Prasad, R., & Nath, A. (2022). Misattribution of hematological malignancies to endemic infectious diseases in rural Asian populations. *Tropical Doctor*, 52(1), 45–51. https://doi.org/10.1177/00494755211048210

Arber, D. A., Orazi, A., Hasserjian, R., Thiele, J., Borowitz, M. J., Le Beau, M. M., Bloomfield, C. D., Cazzola, M., & Vardiman, J. W. (2016). The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*, 127(20), 2391–2405. https://doi.org/10.1182/blood-2016-03-643544

Arora, R. S., Eden, T., & Pizer, B. (2010). The burden of childhood cancer in India: A review of available data. *Pediatric Blood & Cancer*, 54(7), 1033–1037. https://doi.org/10.1002/pbc.22442

Asthana, S., Chauhan, S., & Labani, S. (2017). Cancer incidence and mortality in rural India: Absolute numbers and trends. *Indian Journal of Medical and Paediatric Oncology*, 38(4), 450–457. https://doi.org/10.4103/ijmpo.ijmpo_188_16

Atun, R., Jaffray, D. A., Barton, M. B., Bray, F., Baumann, M., Vikram, B., Mahmood, U., Zubizarreta, E., Yolero, K., & Gospodarowicz, M. (2015). Expanding global access to radiotherapy. *The Lancet Oncology*, 16(10), 1153–1186. https://doi.org/10.1016/S1470-2045(15)00222-3

Bakshi, S., & Basu, S. (2023). Patient navigation programs in public cancer hospitals: Bridging entitlement and actual health coverage. *Journal of Cancer Policy*, 35, Article 100399.

https://doi.org/10.1016/j.jcpo.2023.100399

Balarajan, Y., Selvaraj, S., & Subramanian, S. V. (2011). Healthcare and equity in India. *The Lancet*, 377(9764), 505–515. https://doi.org/10.1016/S0140-6736(10)61894-6

Bates, I., & Maitland, K. (2006). Diagnostic capability in primary healthcare: The neglected foundation of global health systems. *Tropical Medicine & International Health*, 11(4), 411–414. https://doi.org/10.1111/j.1365-3156.2006.01584.x

Bhutani, M., Vora, A., Kumar, L., & Kochupillai, V. (2002). Hematological malignancies in India: An epidemiological review. *Medical Oncology*, 19(3), 139–146. [Corrected journal name from JAPI to Medical Oncology]

Bourpetruc, S., Deak, H., Chinen, K., & Patel, V. (2020). Diagnostic gaps in primary healthcare in rural South Asia. *Tropical Medicine & International Health*, 25(7), 812–824. https://doi.org/10.1111/tmi.13398

Briggs, C., Culp, N., Davis, B., d'Onofrio, G., Zini, G., & Machin, S. J. (2020). International Council for Standardization in Haematology (ICSH) recommendations for workplace evaluation of automated digital cell morphology systems. *International Journal of Laboratory Hematology*, 42(1), 12–28. https://doi.org/10.1111/ijlh.13141

Broom, A., Doron, A., & Tovey, P. (2009). The asymmetry of biomedical pluralism: Oncology and traditional healing in India. *Social Science & Medicine*, 69(5), 700–706. https://doi.org/10.1016/j.socscimed.2009.06.020

Central Bureau of Health Intelligence. (2021). *National Health Profile 2021*. Ministry of Health and Family Welfare, Government of India.

Chandy, M. (2008). Stem cell transplantation in developing countries. *Bone Marrow Transplantation*, 42(Suppl 1), S81–S83. https://doi.org/10.1038/bmt.2008.121

Chandy, M., & Boopathy, V. (2015). Development of clinical hematology services in India: Historical context and future imperatives. *Indian Journal of Hematology and Blood Transfusion*, 31(3), 301–307. https://doi.org/10.1007/s12288-015-0551-x

Chaudhuri, S. (2005). *The pharmaceutical industry in India: A public health perspective*. Oxford University Press.

Chaudhuri, S., Sarkar, M., & Mukherjee, A. (2024). Comparative evaluation of AI-driven peripheral blood smear digitization against manual microscopy in detecting acute blast populations. *Indian Journal of Pathology and Microbiology*, 67(2), 289–295. https://doi.org/10.4103/ijpm.ijpm_412_23

Damani, A., Ghoshal, S., & Salins, N. (2024). Financial toxicity in cancer palliative care in India: Addressing existence and beyond. *ecancermedicalscience*, 18, Article 1820. https://doi.org/10.3332/ecancer.2024.1820

Das, J., Holla, A., Mohpal, A., & Muralidharan, K. (2016). Quality and accountability in healthcare delivery: Audit study of primary

care in rural and urban India. *American Economic Review*, 106(12), 3738–3773. https://doi.org/10.1257/aer.20150499

Dastidar, T. R., Roul, P., Sahu, A., & Kumar, P. (2023). Validation of an automated digital cell morphology analyzer (SigTuple AI100) for peripheral blood smear differential counts. *International Journal of Laboratory Hematology*, 45(5), 690–698. https://doi.org/10.1111/ijlh.14088

Datta, S. S., Ghose, S., Mukherjee, S., & Ray, S. (2022). Staffing gaps in district cancer care facilities across Tier-2 and Tier-3 cities in Eastern India. *Journal of Cancer Policy*, 31, Article 100318. https://doi.org/10.1016/j.jcpo.2021.100318

Dharmshaktu, P., Jaiswal, S. R., Chakraborty, S., & Chakrabarti, S. (2021). Stem cell transplantation in India: Current status, challenges, and regional inequities. *Bone Marrow Transplantation*, 56(8), 1790–1798. https://doi.org/10.1038/s41409-021-01310-8

Dikshit, R., Gupta, P. C., Ramasundarahettige, C., Gajalakshmi, V., Aleksandrowicz, L., Badwe, R., Kumar, R., Roy, S., Suraweera, W., & Jha, P. (2012). Cancer mortality in India: A nationally representative survey. *The Lancet*, 379(9828), 1807–1816. https://doi.org/10.1016/S0140-6736(12)60358-4

Engel, N., Wachter, K., Pai, M., & Gallarda, J. (2019). Private sector diagnostic landscape in India: Implications for healthcare equity. *BMC Health Services Research*, 19(1), Article 451. https://doi.org/10.1186/s12913-019-4279-0

Engelgau, M. M., Karan, A., & Mahal, A. (2012). The economic impact of non-communicable diseases on households in India. *Globalization and Health*, 8(1), Article 9. https://doi.org/10.1186/1744-8603-8-9

Etcheverry, A., Arboleya, S., & Lopez, G. (2023). Cloud-connected digital tele-pathology models for rural diagnostic empowerment: A systemic evaluation. *Telemedicine and e-Health*, 29(8), 1140–1151. https://doi.org/10.1089/tmj.2022.0341

Ganesan, P., Sagar, T. G., Dubashi, B., Radhakrishnan, V., Rajendranath, R., Joseph, R., & Tenali, S. G. (2018). Safe delivery of outpatient maintenance therapy for acute lymphoblastic leukemia in regional centers. *Pediatric Blood & Cancer*, 65(10), Article e27290. https://doi.org/10.1002/pbc.27290

Garg, C. C., & Karan, A. K. (2009). Reducing out-of-pocket expenditures and improving equity in financial risk protection in India. *Bulletin of the World Health Organization*, 87(3), 199–206. https://doi.org/10.2471/blt.08.051961

Gautham, M., Kishora, N., Shyamprasad, S., & Bloom, G. (2014). Informal healthcare providers in rural India: Potential allies or persistent risks in cancer early detection? *Health Policy and Planning*, 29(Suppl 1), i29–i38. https://doi.org/10.1093/heapol/czu064

George, I. A., Bhattacharjee, A., & Roy, S. (2021). Non-medical out-of-pocket costs in cancer treatment: Travel, accommodation, and caregiver

income loss in rural Eastern India. *Supportive Care in Cancer*, 29(9), 5105–5112. https://doi.org/10.1007/s00520-021-06078-4

Gogia, A., Sharma, A., Raina, V., Kumar, L., Vishnubhatla, S., & Sharma, M. C. (2018). Assessment of clinico-pathological profile of acute leukemias in a tertiary care institute in Northern India. *Indian Journal of Cancer*, 55(2), 145–149. https://doi.org/10.4103/ijc.IJC_482_17

Goss, P. E., Strasser-Weippl, K., Lee-Bychkovsky, B. L., Fan, L., Li, J., Chavarri-Guerra, Y., Liedke, P. E., Pramesh, C. S., Badwe, R. A., & Chan, A. (2014). Challenges to effective cancer control in China, India, and Russia. *The Lancet Oncology*, 15(5), 489–538. https://doi.org/10.1016/S1470-2045(14)70029-4

Government of Karnataka Health Department. (2025). *Notification on decentralized daycare chemotherapy infrastructure in 31 district hospitals*. State Health Secretariat. [Official government document—acceptable as grey literature]

Government of Maharashtra. (2025). *Establishment of first public bone marrow transplant facility at CPR Hospital Kolhapur under AB PM-JAY*. Department of Medical Education and Drugs. [Official government document—acceptable as grey literature]

Gratwohl, A., Pasquini, M. C., Aljurf, M., Atsuta, Y., Baldomero, H., Foeken, L., Gratwohl, M., Bouzas, L. F., Confer, D., & Frauendorfer, K. (2015). One million haemopoietic stem-cell transplants: A retrospective observational study. *The Lancet Haematology*, 2(3), e91–e100. https://doi.org/10.1016/S2352-3026(15)00028-9

Gupta, A., & Parikh, P. (2022). Generic targeted agents in hematological malignancies: Transforming survival in low- and middle-income nations. *Current Hematologic Malignancy Reports*, 17(3), 145–154. https://doi.org/10.1007/s11899-022-00668-1

Gupta, R., Rahman, K., Verma, S. P., & Nath, U. K. (2020). Cytogenetic profile of adult acute myeloid leukemia in India and its impact on treatment response. *Annals of Hematology*, 99(6), 1245–1254. https://doi.org/10.1007/s00277-020-04022-w

Gupta, S., Howard, S. C., Hunger, S. P., Antillon, F. G., Metzger, M. L., Israels, T., Harif, M., & Magrath, I. (2015). Treating childhood acute lymphoblastic leukemia in low- and middle-income countries. *Pediatric Blood & Cancer*, 62(3), 387–393. https://doi.org/10.1002/pbc.25251

Gupta, S., Yeh, S., Martina, C., & Banavali, S. (2022). Financial navigation and holistic psychosocial support reduce abandonment in pediatric leukemias in resource-limited settings. *JCO Global Oncology*, 8, Article e2100340. https://doi.org/10.1200/GO.21.00340

Halder, R. (2025). Structural disparities and workforce maldistribution in Indian hematology care. *Blood*, 146(Suppl 1), Article A4102. https://doi.org/10.1182/blood-2025-146-A4102 [Author must verify this is a full paper, not just a conference abstract]

Horton, R. (2015). Offline: What is cancer care for poor people? *The Lancet*, 386(10011), 2376. https://doi.org/10.1016/S0140-6736(15)01188-6

Horton, S., Gauvreau, C. L., Gelband, H., Jha, P., & Sankaranarayanan, R. (2015). *Disease control priorities: Volume 3. Cancer* (3rd ed.). World Bank.

Howard, S. C., Metzger, M. L., Wilimas, J. A., Quintana, Y., Pui, C. H., Robison, L. L., & Ribeiro, R. C. (2008). Childhood cancer cure in poor countries: Total therapy protocols for acute lymphoblastic leukemia in low-income countries. *British Journal of Haematology*, 140(4), 457–466. https://doi.org/10.1111/j.1365-2141.2007.06948.x

Hudecek, M., & Ivics, Z. (2023). CAR-T cell therapy goes affordable: Lessons from indigenous manufacturing in India. *Cell Stem Cell*, 30(11), 1401–1403. https://doi.org/10.1016/j.stem.2023.10.005

India State-Level Disease Burden Initiative Cancer Collaborators. (2018). The burden of cancers and their variations across the states of India: The Global Burden of Disease Study 1990–2016. *The Lancet Oncology*, 19(10), 1289–1306. https://doi.org/10.1016/S1470-2045(18)30447-9

Indian Council of Medical Research. (2023). *Call for action: Strengthening medical oncology human resources in India* (Special Report). ICMR. [Author must verify this document is publicly available online]

Jaiswal, S. R., & Chakrabarti, S. (2016). Allogeneic hematopoietic stem cell transplantation in resource-limited settings: Challenges and solutions from India. *Current Hematologic Malignancy Reports*, 11(5), 385–393. https://doi.org/10.1007/s11899-016-0338-0

Jha, P., Gajalakshmi, V., Gupta, P. C., Kumar, R., Peto, R., Yong, S. N., Choi, B. C., & Ranson, M. K. (2006). Prospective study of one million deaths in India: Background, design, and methods. *Results in Premature Mortality*, 15(1), 20–29. https://doi.org/10.1016/j.annepidem.2005.02.009

Joshi, R., Alim, M., Kengne, A. P., Jan, S., Maulik, P. K., Peiris, D., & Patel, A. A. (2014). Task-shifting for non-communicable disease management in low- and middle-income countries: A systematic review. *PLoS ONE*, 9(8), Article e103754. https://doi.org/10.1371/journal.pone.0103754

Kakkar, F., Radhakrishnan, V., & Banavali, S. (2019). Capacity constraints and wait-list mortality in public stem cell transplant units in India. *Journal of Global Oncology*, 5, 1–9. https://doi.org/10.1200/JGO.19.00112

Kampuanais, P., Treeprasertsuk, S., & Natesuwan, S. (2022). Artificial intelligence in hematopathology: Digital morphology analysis in low-resource settings. *Acta Haematologica*, 145(4), 380–392. https://doi.org/10.1159/000522105

Kastor, A., & Mohanty, S. K. (2018). Disease-specific out-of-pocket health expenditure and catastrophic spending in India. *BMC Health Services Research*, 18(1), Article 603. https://doi.org/10.1186/s12913-018-3412-8

Keshri, V. R., & Jagnoor, J. (2022). Capacity of primary health care frontline workers for early cancer symptom identification: A systematic evaluation in rural Bihar. *Journal of Family Medicine and Primary Care*, 11(6), 2489–2495. https://doi.org/10.4103/jfmpc.jfmpc_2011_21

Krishnan, S., Sharma, S., & Mukherjee, S. (2021). Training frontline community health workers (ASHAs) for early cancer symptom navigation in rural India. *BMC Public Health*, 21(1), Article 1142. https://doi.org/10.1186/s12889-021-11180-2

Kumar, L., Boya, R. R., Shah, R., Sharma, A., & Aggarwal, S. (2021). Multiple myeloma: Indian data and perspective. *Indian Journal of Medical Research*, 154(2), 210–222. https://doi.org/10.4103/ijmr.IJMR_1120_20 [Author must verify the article title and content match the citation claim]

Kumar, R., Singh, S., Mahto, S. K., & Sharma, P. (2026). Clinico-epidemiological profile of hematological malignancies presenting at a tertiary care center in Eastern India: A 2-year cross-sectional study. *Indian Journal of Medical and Paediatric Oncology*, 47(1), 34–41. https://doi.org/10.1055/s-0045-1782301

Lall, O. O., Kazi, M., & Parihar, M. (2023). Hematopathology workforce shortages in South Asia: Challenges and structural solutions. *Journal of Clinical Pathology*, 76(4), 215–221. https://doi.org/10.1136/jcp-2022-208450

Lambert, H., & Wood, K. (2005). Therapeutic pathways and delays in care-seeking for chronic symptoms in rural central India. *Social Science & Medicine*, 61(1), 243–255. https://doi.org/10.1016/j.socscimed.2004.11.042

Magrath, I., Steliarova-Foucher, E., Epelman, S., Ribeiro, R. C., Harif, M., Li, C. K., Eden, T., Hesseling, P., Howard, S. C., & Howard, R. (2013). Paediatric cancer in low-income and middle-income countries. *The Lancet Oncology*, 14(3), e104–e116. https://doi.org/10.1016/S1470-2045(13)70008-1

Malhotra, H., Radhakrishnan, V., Rego, E. M., & Almeida, A. (2020). Chronic myeloid leukemia care in low- and middle-income countries. *JCO Global Oncology*, 6, 1044–1052. https://doi.org/10.1200/GO.20.00048

Malhotra, H., & Sharma, M. (2024). Tele-hematology in India: Bridging the urban-rural diagnostic and therapeutic divide. *Current Opinion in Hematology*, 31(2), 95–102. https://doi.org/10.1097/MOH.0000000000000798

Mallath, M. K., Taylor, D. G., Badwe, R. A., Rath, G. K., Shanta, V., Pramesh, C. S., Digumarti, R., Sebastian, P., Sullivan, R., & Arney, S. (2014). The growing burden of cancer in India: Epidemiology and social context. *The Lancet Oncology*, 15(6), e205–e212. https://doi.org/10.1016/S1470-2045(14)70115-9

Mathew, A. (2018). Global survey of clinical oncology workforce: India country analysis. *Journal of Global Oncology*, 4, 1–12. https://doi.org/10.1200/JGO.17.00188

Mathur, P., Sathishkumar, K., Chaturvedi, M., Das, P., Sudarshan, K. L., Santhappan, S., Nallasivam, A., Vijay, C. R., & Prashant, M. (2020). Cancer statistics, 2020: Report from National Cancer Registry Programme, India. *JCO Global Oncology*, 6, 1063–1075. https://doi.org/10.1200/GO.20.00122

Ministry of Finance. (2025). *Union budget 2025–26: Health sector allocations and establishment of 200 district day care cancer centres*. Government of India. [Official government document—acceptable as grey literature]

Ministry of Health and Family Welfare. (2017). *Operational guidelines for prevention, screening and control of common non-communicable diseases*. Government of India. [Official government document—acceptable as grey literature]

Ministry of Health and Family Welfare. (2023). *eSanjeevani national telemedicine service: Crossing 100 million tele-consultations milestone*. Government of India Press Release. [Official government document—acceptable as grey literature]

Mishra, F., Singh, P., & Karia, S. (2022). Tele-oncology frameworks for remote supportive care and chemotherapy supervision in underserved India. *Journal of Telemedicine and Telecare*, 28(6), 415–423. https://doi.org/10.1177/1357633X20952110

Mishra, K., Basu, S., & Rath, B. (2019). Hematological manifestations of systemic infectious diseases mimicking acute leukemias in rural Indian clinics. *Journal of Tropical Pediatrics*, 65(4), 356–363. https://doi.org/10.1093/tropej/fmy061

Mohanti, B. K., Mukhopadhyay, A., Das, S., Sharma, K., & Dash, S. (2011). Economic vulnerability and financial catastrophe in cancer care: A prospective multi-center trial in rural India. *Journal of Cancer Research and Therapeutics*, 7(2), 142–148. https://doi.org/10.4103/0973-1482.82918

Mostert, S., Arora, R. S., Arreola, M., Bagai, P., Friedrich, P., Ribeiro, R. C., & Kaspers, G. J. (2011). Abandonment of treatment for childhood cancer: A position statement of a SIOP PODC Working Group. *The Lancet Oncology*, 12(8), 719–720. https://doi.org/10.1016/S1470-2045(11)70128-0

Nair, R., Arora, A., Viswabandya, A., & Chandy, M. (2019). Epidemiology and treatment outcomes of multiple myeloma in rural western India: A community registry analysis. *Leukemia Research*, 80, 1–7. [Corrected journal name from Leukemia & Lymphoma to Leukemia Research; author must verify exact volume/pages]

Narula, G., Prasad, M., Banavali, S., Kurkure, P., & Gupta, S. (2017). ImPaCCT (Improving Paediatric Cancer Care and Treatment) Foundation: A holistic support model mitigating abandonment in childhood cancer. *Indian Journal of Cancer*, 54(1), 112–118. https://doi.org/10.4103/ijc.IJC_142_17

National Centre for Disease Informatics and Research. (2021). *Report on hospital based cancer registries 2021*. Indian Council of Medical Research.

National Health Authority. (2024). *Ayushman Bharat PM-JAY: Annual performance report 2023-24*. Government of India.

National Pharmaceutical Pricing Authority. (2019). *Price fixation and cap strategy for 42 essential non-scheduled cancer drugs*. Ministry of Chemicals and Fertilizers, Government of India.

Neelapu, S. S., Tummala, S., Kebriaei, P., Wierda, W., Gutierrez, C., Locke, F. L., Komanduri, K. V., Lin, Y., Jain, N., & Daver, N. (2018). Chimeric antigen receptor T-cell therapy—assessment and management of toxicities. *Nature Reviews Clinical Oncology*, 15(1), 47–62. https://doi.org/10.1038/nrclinonc.2017.148

Overcoming Prices of Targeted Oncology Therapy. (2016). Global impact of generic imatinib availability. *New England Journal of Medicine*, 374(19), 1801–1804. https://doi.org/10.1056/NEJMp1601230

Pai, N. P., Vadnais, C., Denkinger, C., Engel, N., & Pai, M. (2012). Point-of-care diagnostics for global health: Biosensors and digital health integration. *PLoS Medicine*, 9(9), Article e1001301. https://doi.org/10.1371/journal.pmed.1001301

Patkar, N., Subramanian, P. G., Tembhare, P., Chatterjee, G., & Gujral, S. (2017). Flow cytometric immunophenotyping in acute leukemias: Practical consensus recommendations for Indian laboratories. *Indian Journal of Hematology and Blood Transfusion*, 33(2), 155–164. https://doi.org/10.1007/s12288-017-0808-1

Paul, V. K., Patel, V., Jha, P., Laxminarayan, R., Reddy, K. S., & Hotez, P. J. (2011). Universal health coverage for India: Accelerating progress. *The Lancet*, 377(9766), 668–679. https://doi.org/10.1016/S0140-6736(10)62160-5

Pawaskar, A., Kulkarni, S., Patil, A., & Deshmukh, R. (2025). Scalable tertiary-rural partnership for delivery of advanced leukemia therapy and stem cell transplant in tribal Maharashtra: A retrospective cohort of 60 patients. *The Lancet Regional Health - Southeast Asia*, 32, Article 100412. https://doi.org/10.1016/j.lansea.2024.100412

Peters, D. H., Yazbeck, A. S., Sharma, R. R., Ramana, G. N. V., Pritchett, L. H., & Wagstaff, A. (2002). *Better health systems for India's poor: Findings, analysis, and options*. World Bank Publications.

Pramesh, C. S., Badwe, R. A., Bhoo-Pathy, N., Sullivan, R., Lambertini, M., & Booth, C. M. (2022). Priorities for cancer research in low-income and middle-income countries. *The Lancet Oncology*, 23(4), e171–e180. https://doi.org/10.1016/S1470-2045(21)00742-X

Pramesh, C. S., Badwe, R. A., & Sinha, R. (2014). The National Cancer Grid of India: A collaborative network for uniform standards of care. *The Lancet Oncology*, 15(8), e340–e343. https://doi.org/10.1016/S1470-2045(14)70163-9

Pramesh, C. S., Kapoor, R., & Gupta, S. (2023). Decentralized chemotherapy delivery models in Tier-2 and Tier-3 Indian districts: Clinical safety and economic

impact. *Bulletin of the World Health Organization*, 101(9), 980–991. [Corrected journal name from The Lancet Oncology to Bull WHO; author must verify exact volume/pages]

Press Information Bureau. (2025). *Towards a cancer-free India: Progress report on National Programme for Prevention and Control of Cancer, Diabetes, Cardiovascular Diseases and Stroke (NPCDCS)*. Ministry of Health and Family Welfare. [Official government document—acceptable as grey literature]

Prinja, S., Bahuguna, P., Gupta, I., Chowdhury, S., & Trivedi, M. (2021). Assessment of financial protection under Ayushman Bharat PM-JAY in North India. *PLoS ONE*, 16(8), Article e0255915. https://doi.org/10.1371/journal.pone.0255915

Purwar, R., Narula, G., Joshi, K., & Hudecek, M. (2023). First-in-human phase I/II clinical trial of indigenous CD19 CAR-T cells (NexCAR19) in relapsed/refractory B-cell malignancies in India. *Blood*, 142(Suppl 1), Article A102. https://doi.org/10.1182/blood-2023-142-A102

Radhakrishnan, V., Dhanushkodi, E., Ganesan, P., Rajendranath, R., Ganesan, T. S., & Sagar, T. G. (2021). Outcome of acute lymphoblastic leukemia in children and young adults: A single-center experience from India. *Pediatric Blood & Cancer*, 68(5), Article e28912. https://doi.org/10.1002/pbc.28912

Rajappa, S., Joshi, A., Biswas, G., & Vaid, A. (2020). Impact of generic imatinib on CML outcomes in India: A 15-year real-world cohort analysis. *South Asian Journal of Cancer*, 9(3), 142–147. https://doi.org/10.4103/sajc.sajc_42_20

Rajkumar, S. V. (2022). Multiple myeloma: 2022 update on diagnosis, risk-stratification, and management. *American Journal of Hematology*, 97(8), 1086–1107. https://doi.org/10.1002/ajh.26590

Ramamoorthy, T., Sarveswaran, G., Nath, A., Vijayakumar, S., & Mathur, P. (2025). Distance travelled for cancer care and its association with stage and treatment patterns: Analysis from hospital-based cancer registries in India. *Indian Journal of Medical Research*, 161(2), 180–189. https://doi.org/10.4103/ijmr.ijmr_892_24

Rao, K. D., Shahrawat, R., & Bhatnagar, A. (2016). Composition and distribution of the health workforce in India: Estimates based on certified data. *Human Resources for Health*, 14(1), Article 30. https://doi.org/10.1186/s12960-016-0126-7

Roy, S., Ghosh, A., & Chatterji, M. (2024). Infrastructure requirements for cellular therapy delivery in low- and middle-income countries. *Cytotherapy*, 26(2), 115–122. https://doi.org/10.1016/j.jcyt.2023.10.004

Roy, S., Sharma, A., & Kumar, L. (2019). Biosimilar rituximab in the management of Non-Hodgkin Lymphoma: Real-world efficacy, safety, and pharmacoeconomic data from India. *Leukemia & Lymphoma*, 60(11), 2712–2720. https://doi.org/10.1080/10428194.2019.1605072

Sengar, M., Jain, H., Murthy, V., & Gupta, S. (2021). Impact of decentralized care on treatment adherence and

financial expenditure among adult lymphoma patients in India. *JCO Global Oncology*, 7, 1210–1218. https://doi.org/10.1200/GO.21.00140

Sharma, K., Zodpey, S., Preetish, A., & Quazi, S. Z. (2020). Specialization choice and rural service retention among Indian medical postgraduates: A cross-sectional survey. *Health Policy and Planning*, 35(5), 580–589. https://doi.org/10.1093/heapol/czaa012

Sharma, L., Purohit, A., Vyas, S., & Singh, R. (2024). Economic burden and out-of-pocket expenditure among families of patients with hematological malignancies in Western Rajasthan. *Indian Journal of Medical and Paediatric Oncology*, 45(3), 210–218. https://doi.org/10.1055/s-0044-1779012

Sharma, S., Mishra, A. K., & Singh, S. (2021). Diagnostic delays and misdiagnoses of lymphadenopathy in rural primary care: Tuberculosis versus lymphoma. *Journal of Family Medicine and Primary Care*, 10(3), 1120–1126. https://doi.org/10.4103/jfmpc.jfmpc_1890_20

Sheikh, K., Rajkumari, B., Jain, K., & Rao, K. D. (2012). What drives health workforce retention in rural underserved districts? Qualitative insights from Indian public sector doctors. *BMJ Open*, 2(4), Article e001259. https://doi.org/10.1136/bmjopen-2012-001259

Subramanian, S. V., Bhatia, M., & Kawachi, I. (2008). Disparities in health and healthcare access in India: A persistent challenge. *Health Affairs*, 27(4), 979–990. https://doi.org/10.1377/hlthaff.27.4.979

Sullivan, R., Kowalychin, J. P., Sarfati, D., Tremblay, A., & Aggarwal, A. (2015). Global cancer challenges and crisis: Equitable health-system responses. *The Lancet Oncology*, 16(11), 1193–1206. https://doi.org/10.1016/S1470-2045(15)00084-4

Swaminathan, R., Rama, R., & Shanta, V. (2008). Lack of population-based cancer registries in rural India: Impact on health planning and resource allocation. *Bulletin of the World Health Organization*, 86(11), 880–886. https://doi.org/10.2471/blt.07.047860

Swaminathan, S., Qureshi, A., & Vijayaraghavan, K. (2023). Community health worker-led health literacy and adherence monitoring in pediatric leukemia care in rural South India. *The Lancet Global Health*, 11(Suppl 1), S14. https://doi.org/10.1016/S2214-109X(23)00112-8

Swerdlow, S. H., Campo, E., Pileri, S. A., Harris, N. L., Stein, H., Siebert, R., Advani, R., Ghielmini, M., Salles, G. A., & Zelenetz, A. D. (2016). The 2016 revision of the World Health Organization classification of lymphoid neoplasms. *Blood*, 127(20), 2375–2390. https://doi.org/10.1182/blood-2016-01-643569

Takiar, R., Nadayil, D., & Nandakumar, A. (2010). Projections of number of cancer cases in India (2010-2020) by cancer groups. *Asian Pacific Journal of Cancer Prevention*, 11(4), 1045–1049.

Tembhare, P., Narula, G., Kurkure, P., & Banavali, S. (2019). Factors

predicting treatment refusal and abandonment in pediatric acute lymphoblastic leukemia: Single-center cohort study from Mumbai. *Psycho-Oncology*, 28(4), Article e27581. [Corrected journal name from Pediatric Blood & Cancer to Psycho-Oncology; author must verify exact volume/pages]

Tripathi, S., Sinha, S., & Kapoor, R. (2025). Health-seeking pathways and diagnostic lag times in pediatric acute leukemias in North India. *Journal of Pediatric Hematology/Oncology*, 47(2), 112–119. https://doi.org/10.1097/MPH.0000000000002890 [Author must verify this reference exists as cited]

Udayakumar, M., Mathews, V., George, B., Srivastava, A., & Balasubramanian, P. (2020). Cost-effectiveness analysis of allogeneic stem cell transplantation for acute leukemia in a public sector tertiary hospital in India. *Indian Journal of Hematology and Blood Transfusion*, 36(3), 478–485. https://doi.org/10.1007/s12288-020-01258-2

US Food and Drug Administration. (2023). *510(k) Clearance K230411: SigTuple AI100 Automated Digital Hematology System*. US FDA.

Ved, R., Scott, K., Gupta, G., & George, A. (2019). How are ASHA workers supported to perform their roles? A review of performance interventions in India. *BMC Health Services Research*, 19(1), Article 102. https://doi.org/10.1186/s12913-019-3917-x

Weller, D., Vedsted, P., Rubin, G., Walter, F. M., Emery, J., Scott, S., Campbell, C., Andersen, R. S., Hamilton, W., & Olesen, F. (2012). The European Diagnostic Interval Assessment Grant (EDIAG) group consensus statement on defining and measuring diagnostic intervals in cancer research. *British Journal of Cancer*, 106(2), 286–290. https://doi.org/10.1038/bjc.2011.563

Wilson, M. L., Fleming, K. A., Kuti, M. A., Looi, L. M., Lago, N., & Ru, K. (2018). Access to pathology and laboratory medicine services in low-income and middle-income countries. *The Lancet*, 391(10133), 1927–1938. https://doi.org/10.1016/S0140-6736(18)30458-6

World Health Organization. (2021). *Selection and use of essential in vitro diagnostics: Report of the third meeting of the WHO Strategic Advisory Group of Experts on In Vitro Diagnostics*. World Health Organization.

World Health Organization. (2022). *WHO guidelines for establishing decentralized outpatient chemotherapy services in low-resource settings*. World Health Organization.

Yeole, B. B. (2008). Trends in cancer incidence in female breast and cervix in urban and rural population of India. *Asian Pacific Journal of Cancer Prevention*, 9(3), 403–408.

Zafar, S. Y., & Abernethy, A. P. (2013). Financial toxicity: An emerging side effect of high-impact cancer care. *Translational Behavioral Medicine*, 3(4), 433–442. https://doi.org/10.1007/s13142-013-0215-3

Journal of Rural Advancement
[ISSN 2347-2561 (P); 2583-6102 (E)]
Vol. 12, Issue 02, October 2024

Apoptosis Dysregulation in Lung Cancer: Therapeutic Resistance Mechanisms and Implications for Rural Healthcare Advancement in India

Tushar Sharma

Department of Biotechnology and Bioinformatics, North-Eastern Hill University, Shillong 793022, Meghalaya, India

Email: tusharkisho@gmail.com

Received on: May 18, 2024; Accepted on: June 12, 2024

Abstract

Apoptosis is a fundamental mechanism for maintaining cellular homeostasis and eliminating damaged cells. In lung cancer, this tightly regulated process is frequently disrupted, enabling malignant cells to evade death and persist despite cytotoxic therapy. The two principal apoptotic pathways—intrinsic and extrinsic—are both implicated in this disruption. The intrinsic pathway is governed by the Bcl-2 family of proteins, which control mitochondrial outer membrane permeabilization and downstream caspase activation, while the extrinsic pathway is initiated by death receptors on the cell surface and proceeds through the death-inducing signaling complex. Mutations in key regulators such as TP53, overexpression of anti-apoptotic proteins (notably MCL-1 in non-small-cell lung cancer), and defects in death-receptor signaling collectively raise the apoptotic threshold and underlie resistance to chemotherapy, radiotherapy, and targeted agents. Emerging strategies—BH3 mimetics, MCL-1-selective inhibitors, pharmacological p53 reactivation, TRAIL-receptor agonists, and rational combinations—show promise but remain dependent on functional biomarkers of apoptotic dependence rather than static gene-expression data. This review synthesizes the mechanistic literature on apoptosis dysregulation in lung cancer and connects it to the rural Indian context, where rising incidence, biomass-fuel exposure, late-stage diagnosis, and limited access to specialized oncology require affordable, scalable, and decentralized strategies. We argue that apoptosis-targeted therapy will improve rural outcomes only if paired with upstream risk reduction, decentralized diagnostic pathways including telepathology, low-cost tumor-selective delivery platforms, and validation of functional biomarkers in resource-limited settings.

Keywords: Apoptosis, lung cancer, MCL-1, BH3 mimetics, therapeutic resistance, telepathology, rural health, India.

1. Introduction

Lung cancer is the leading cause of cancer-related mortality worldwide and accounts for a substantial and growing share of the cancer burden in India. Recent Global Burden of Disease analyses indicate that tracheal, bronchus, and lung cancer accounted for 8.73% of all cancer fatalities

in India in 2021, with age-standardized incidence and mortality rates rising steadily since 1990 . Although national cancer statistics have historically been derived from urban population-based registries, hospital-based data and regional audits consistently indicate that a large and increasing proportion of Indian lung cancer patients present from rural and semi-urban districts (Indian Council of Medical Research, 2020). Rural populations face a distinctive risk profile: a high prevalence of smoking and smokeless tobacco use, widespread exposure to biomass fuel combustion in poorly ventilated homes, delayed presentation due to limited screening infrastructure, and restricted access to specialized thoracic oncology and molecular testing. These factors converge to produce later-stage diagnosis and poorer outcomes than those observed in metropolitan centers.

Apoptosis—programmed cell death—is essential for tissue homeostasis and for the elimination of damaged or potentially malignant cells (Elmore, 2007). Evasion of apoptosis is a hallmark of cancer (Hanahan & Weinberg, 2011), and in lung cancer the apoptotic machinery is disabled at multiple nodes: loss of p53 function, overexpression of anti-apoptotic Bcl-2 family proteins, defects in death-receptor signaling, and rewiring of survival kinase pathways. Because conventional chemotherapy and radiotherapy depend on the induction of apoptosis for their cytotoxic effect, these lesions are directly responsible for therapeutic resistance and treatment failure.

This review has two aims. First, it synthesizes the molecular literature on intrinsic and extrinsic apoptosis dysregulation in lung cancer, with attention to the lung-cancer-specific features—particularly the dominant role of MCL-1 in non-small-cell lung cancer (NSCLC)—that distinguish it from other tumor types. Second, it examines how these mechanistic insights can be translated into rural healthcare advancement in India, where the rising burden of lung cancer and the scarcity of tertiary oncology capacity require affordable, scalable, and context-sensitive strategies. The review deliberately connects molecular mechanism to health-system design, arguing that apoptosis-targeted therapy will remain an urban tertiary-care privilege unless it is embedded in a decentralized diagnostic and delivery framework.

2. The Intrinsic Apoptotic Pathway

The intrinsic, or mitochondrial, pathway is regulated by the Bcl-2 family, which comprises both pro-apoptotic members (Bax, Bak, and the BH3-only proteins BIM, PUMA, BID, and others) and anti-apoptotic members (Bcl-2, Bcl-xL, MCL-1, Bcl-w, and A1). In response to cellular stress, BH3-only proteins activate Bax and Bak, promoting mitochondrial outer membrane permeabilization and the release of cytochrome c into the cytosol (Green & Kroemer, 2004; Youle & Strasser, 2008). Cytochrome c associates with Apaf-1 and procaspase-9 to form the apoptosome, which activates caspase-9 and, in turn, effector caspases such as caspase-3 (Acehan et al., 2002; Riedl & Salvesen, 2007; Bratton & Salvesen, 2010). Anti-apoptotic proteins preserve mitochondrial integrity by sequestering Bax and Bak or by binding BH3-only proteins, and the balance between these opposing factions determines cell fate (Cory & Adams, 2002; Willis et al., 2005).

The regulation of this pathway is layered. BH3-only proteins act as stress sensors that either directly activate Bax and Bak or antagonize their anti-apoptotic sequesters

(Ren et al., 2010; Moldoveanu et al., 2013). Post-translational modifications—phosphorylation, ubiquitination, and proteolytic cleavage—further modulate the activity and stability of Bcl-2 family members and shift the apoptotic threshold (Deng et al., 2007; Kale et al., 2018). p53 sits upstream of the pathway, transcriptionally inducing Bax and PUMA in response to DNA damage (Vousden & Prives, 2009; Aubrey et al., 2016).

2.1 MCL-1 as the Dominant Anti-Apoptotic Dependency in NSCLC

The identity of the dominant anti-apoptotic protein differs by tumor type, and this has direct therapeutic consequences. In NSCLC, **MCL-1**—not BCL-2—is the principal apoptotic block, and MCL-1 amplification is a common mechanism of resistance to both chemotherapy and BCL-2-selective BH3 mimetics (Willis et al., 2005; Kale et al., 2018). MCL-1 is a pro-survival member of the BCL-2 family that binds with high affinity to the BH3 domains of pro-apoptotic effectors, sequestering them and preventing mitochondrial outer membrane permeabilization . Its short half-life and rapid inducible expression make it a dynamic and adaptive resistance node: cytotoxic therapy that damages DNA can paradoxically upregulate MCL-1 through stress-responsive signaling, creating a survival feedback loop that limits drug efficacy.

Preclinical work in NSCLC models has confirmed that MCL-1 dependence is not merely a correlate of resistance but a causal driver. Dynamic BH3 profiling studies in ALK-rearranged NSCLC have shown that MCL-1 plays a key role in tumor resistance to ALK inhibitors, and that BH3 mimetics targeting MCL-1 can restore sensitivity to these targeted agents by preventing tumor adaptation . This finding has direct clinical implications: ALK inhibitors, while highly effective initially, produce responses that are rarely durable, and the emergence of MCL-1-dependent resistance represents a rational target for combination strategies.

Small-cell lung cancer (SCLC), by contrast, is characterized by near-universal inactivation of TP53 and RB1 and a high apoptotic priming state that contributes to its initial chemosensitivity and subsequent rapid relapse. The distinction between NSCLC (MCL-1-dependent) and SCLC (highly primed, p53/RB1-null) is therefore not academic: it determines which apoptotic dependencies are pharmacologically tractable in each histology.

2.2 Driver-Mutation Context and Apoptotic Priming

Driver-mutation context further shapes apoptotic dependency. EGFR-mutant NSCLC depends on BIM-mediated apoptotic priming for tyrosine kinase inhibitor (TKI)-induced cell death, and BIM deletion polymorphism is a documented cause of TKI resistance. In KRAS-mutant NSCLC—a genotype for which targeted therapy has historically been limited—resistance to emerging KRAS-G12C inhibitors frequently involves MCL-1 or BCL-xL upregulation, and combination strategies that co-target these anti-apoptotic proteins are under active investigation . A review of apoptosis in lung cancer that discusses Bcl-2 and Bcl-xL without addressing MCL-1 dependence and histology-specific priming therefore omits the central determinants of therapeutic response.

3. The Extrinsic Apoptotic Pathway

The extrinsic pathway is initiated by death ligands—FasL and TRAIL—binding to death receptors Fas (CD95/APO-1) and

DR4/DR5. Ligand binding promotes receptor trimerization, recruitment of adaptor proteins, formation of the death-inducing signaling complex (DISC), and activation of procaspase-8 (Ashkenazi & Dixit, 1998; Kischkel et al., 1995; Peter & Krammer, 2003). Active caspase-8 directly activates effector caspases, and in so-called type II cells it also cleaves the BH3-only protein BID, engaging the mitochondrial amplification loop. The pathway is modulated by decoy receptors and by c-FLIP, which inhibits caspase-8 activation at the DISC (Ashkenazi, 2002; Micheau & Tschopp, 2003; Safa, 2012).

In lung cancer, extrinsic pathway defects contribute to immune evasion and therapeutic resistance. Downregulation of Fas and mutations in DR4/DR5 reduce sensitivity to ligand-induced apoptosis, while c-FLIP overexpression blunts DISC-mediated caspase-8 activation (Safa, 2012; Djerbi et al., 1999; Gonzalvez & Ashkenazi, 2010). These alterations are not merely descriptive: they predict failure of TRAIL-receptor agonists and contribute to resistance to immune-mediated killing.

Therapeutic strategies targeting the extrinsic pathway include recombinant TRAIL and agonistic antibodies against DR4/DR5, which preferentially induce apoptosis in malignant cells while sparing normal tissue (Ashkenazi, 2008; Herbst et al., 2010). Clinical activity as monotherapy has been limited by c-FLIP upregulation and death-receptor loss, motivating combination approaches with chemotherapy or Bcl-2-family inhibitors (Dimberg et al., 2013).

3.1 Next-Generation TRAIL Delivery: Mesenchymal Stromal Cell Vehicles

A limitation of recombinant TRAIL is its short half-life and poor tumor penetration. To address this, mesenchymal stromal cells (MSCs) engineered to express TRAIL have been developed as cell-based delivery vehicles. MSCs migrate to tumor sites and can deliver TRAIL locally, overcoming the pharmacokinetic limitations of the soluble protein. Preclinical studies demonstrated that MSC-TRAIL has stronger cytotoxic effects than recombinant TRAIL and synergizes with chemotherapy by engaging both extrinsic and intrinsic pathways simultaneously . A first-in-human phase I trial of umbilical cord-derived MSC-TRAIL (UC-MSCTRAIL) in combination with chemotherapy and immunotherapy in advanced NSCLC has been conducted, representing a translational advance in extrinsic-pathway targeting . While still early-phase, this approach illustrates the principle that effective extrinsic-pathway activation requires engagement of the mitochondrial amplification loop—a principle directly relevant to the combination strategies discussed in Section 6.

4. Dysregulation of Apoptotic Pathways in Lung Cancer

Dysregulation in lung cancer is best understood as a set of convergent lesions that together raise the apoptotic threshold. TP53 mutations—present in the majority of SCLC and a large fraction of NSCLC—impair DNA-damage-induced apoptosis and permit survival of genomically unstable cells (Vousden & Lane, 2007). Constitutive activation of PI3K/AKT and NF-κB signaling provides survival input that further shifts the balance toward persistence (Vivanco & Sawyers, 2002; Hanahan & Weinberg, 2011). On the intrinsic axis, overexpression of anti-apoptotic Bcl-2 family members—with MCL-1 predominating in NSCLC—

sequesters pro-apoptotic effectors and blunts the response to cytotoxic agents.

On the extrinsic axis, reduced Fas and DR4/DR5 expression, DR4/DR5 mutation, and c-FLIP overexpression impair DISC formation and caspase-8 activation, contributing to immune evasion and resistance to death-receptor agonists (Safa, 2012; Djerbi et al., 1999). Importantly, these lesions are often redundant: a tumor may simultaneously overexpress MCL-1, lose p53 function, and downregulate death receptors, so that single-agent targeting of any one node is insufficient.

5. Apoptotic Dysfunction and Therapeutic Resistance

Conventional chemotherapy and radiotherapy depend on apoptosis for cytotoxicity, and resistance to these modalities is frequently explained by the lesions described above. Overexpression of Bcl-2, Bcl-xL, or MCL-1 prevents mitochondrial outer membrane permeabilization and blunts the response to genotoxic agents (Gross et al., 1999; Chen et al., 2005; Kale et al., 2018). TP53 mutations impair transcriptional induction of Bax and PUMA, allowing damaged cells to survive (Vousden & Lane, 2007). Dysregulated PI3K/AKT and NF-κB signaling reinforces survival (Vivanco & Sawyers, 2002; Hanahan & Weinberg, 2011). In the extrinsic pathway, death-receptor loss, TRAIL-receptor mutation, and c-FLIP overexpression all contribute to resistance (Safa, 2012; Djerbi et al., 1999).

Because these mechanisms coexist, single-agent therapies are generally insufficient. This recognition has shifted the field toward combination strategies that lower the apoptotic threshold before or alongside a cytotoxic insult.

6. Therapeutic Strategies Targeting Apoptotic Pathways

6.1 Bcl-2 Family Inhibition: From Pan-BCL-2 to MCL-1 Selectivity

BH3 mimetics—small molecules that mimic BH3-only proteins—bind and inhibit anti-apoptotic Bcl-2 family members, releasing pro-apoptotic effectors and inducing apoptosis. Venetoclax (ABT-199), a BCL-2-selective inhibitor, is approved in chronic lymphocytic leukemia and acute myeloid leukemia (Souers et al., 2013; Roberts et al., 2016). In lung cancer, however, single-agent activity has been limited, in large part because **MCL-1, not BCL-2, is the dominant dependency in NSCLC**.

This recognition has driven the development of MCL-1-selective inhibitors. S63845 binds with high affinity to the BH3-binding groove of MCL-1 and kills MCL-1-dependent cancer cells by activating the BAX/BAK-dependent mitochondrial apoptosis pathway . AZD5991, a macrocyclic small molecule, is highly potent and selective for MCL-1, with demonstrated activity in hematological malignancies and sensitivity in subsets of solid tumors including NSCLC . AMG 176 utilizes a distinct chemotype with long residence times and has demonstrated significant tumor regression in xenograft models . A phase I study of the MCL-1 inhibitor PRT1419 has enrolled patients with advanced solid tumors including NSCLC, indicating that clinical translation of MCL-1 inhibition in lung cancer is underway .

An important caveat is on-target cardiac toxicity: MCL-1 is implicated in normal cardiac myocyte function, and early clinical trials of MCL-1 inhibitors have raised concerns about cardiac side effects that may

limit the therapeutic window . This toxicity profile has two implications for the present review. First, MCL-1 inhibitors will likely require tumor-selective delivery or carefully managed dosing schedules. Second, in rural settings where cardiac monitoring infrastructure is limited, the deployment of MCL-1 inhibitors will require particular caution and may favor oral agents with wider safety margins or tumor-targeted delivery platforms.

6.2 Restoring p53 Function

Pharmacological reactivation of mutant p53—through small molecules such as PRIMA-1/APR-246 (eprenetapopt) and related compounds—aims to restore the transcriptional induction of Bax and PUMA and thereby re-prime the intrinsic pathway. This is distinct from gene-replacement approaches, which introduced wild-type TP53 by viral vector in early lung-cancer trials (Roth et al., 1996). Neither strategy has yet produced durable single-agent benefit in lung cancer, but p53 reactivation remains rational in combination with chemotherapy or BH3 mimetics, because it addresses a transcriptional deficit that BH3 mimetics alone cannot overcome.

6.3 TRAIL-Based Therapies and Cellular Delivery

Recombinant TRAIL and agonistic anti-DR4/DR5 antibodies selectively induce apoptosis in malignant cells (Ashkenazi, 2008; Herbst et al., 2010). Clinical efficacy has been constrained by c-FLIP upregulation, death-receptor downregulation, and the failure of extrinsic pathway activation to engage the mitochondrial amplification loop in resistant tumors. Combination with chemotherapy or Bcl-2-family inhibitors is being explored to overcome these blocks (Dimberg et al., 2013). As noted in Section 3.1, MSC-based TRAIL delivery represents a next-generation approach that addresses the pharmacokinetic limitations of soluble TRAIL and has entered first-in-human testing in NSCLC .

6.4 Combination Strategies

The redundancy of apoptotic blocks in lung cancer argues for regimens that engage multiple nodes simultaneously: Bcl-2 or MCL-1 inhibition with chemotherapy, p53 reactivation with BH3 mimetics, and TRAIL agonists with Bcl-2-family inhibitors (Roberts et al., 2016; Dimberg et al., 2013; Roth et al., 1996). The organizing principle is that apoptosis-targeted agents are most effective when used to lower the apoptotic threshold in the context of a second insult, rather than as monotherapy. The finding that MCL-1 targeting enhances CDDP efficacy in solid tumor models—whereas BCL-2 targeting does not—provides direct preclinical support for this principle and underscores the importance of matching the BH3 mimetic to the tumor's actual apoptotic dependency .

7. Crosstalk Between the Intrinsic and Extrinsic Pathways

The two pathways are not independent. Caspase-8 activated at the DISC cleaves BID; truncated BID translocates to mitochondria, where it promotes Bax/Bak activation and engages the intrinsic pathway (Li et al., 1998; Luo et al., 1998). This mitochondrial amplification loop ensures robust commitment to death and has direct therapeutic implications: effective extrinsic-pathway activation depends on an intact mitochondrial response, and tumors with high c-FLIP expression or death-receptor loss remain resistant even when the receptor is engaged. Conversely, intrinsic-pathway blocks—particularly MCL-1 overexpression—can

limit the efficacy of extrinsic agonists. This interdependence explains why single-agent therapies targeting either pathway often fail, and it provides the rationale for combinations that engage both pathways (Dimberg et al., 2013).

8. Biomarkers and Patient Selection

Clinical experience with apoptosis-targeted agents has made clear that patient selection cannot rely on static gene-mutation or expression data. Bcl-2 overexpression is common in lung cancer but does not reliably predict response to Bcl-2 inhibition; TP53 mutation status alone does not identify which patients will benefit from p53 reactivation; and death-receptor expression levels have not reliably predicted TRAIL-agonist activity.

8.1 Functional Biomarkers: BH3 Profiling and Dynamic BH3 Profiling

Functional and dynamic biomarkers address this gap. **BH3 profiling** measures the proximity of a tumor cell to the apoptotic threshold by quantifying mitochondrial depolarization after exposure to BH3-only peptides, and it identifies the specific anti-apoptotic protein on which a tumor depends—information that static expression data cannot provide (Letai, 2008; Deng et al., 2007). **Dynamic BH3 profiling (DBP)** extends this approach by measuring changes in apoptotic priming after ex vivo drug exposure, functioning as a functional precision-medicine assay analogous to an antibiogram . In NSCLC, DBP has been shown to predict response to ALK inhibitors and to identify MCL-1 as a resistance node, supporting its use as a patient-selection tool for apoptosis-targeted combinations . In solid tumor models, DBP has predicted which BH3 mimetic enhances chemotherapy efficacy, with MCL-1 and BCL-xL targeting—but not BCL-2 targeting—showing synergy with cisplatin .

8.2 Circulating Biomarkers

Circulating biomarkers, including cleaved cytokeratin 18 (a caspase-cleavage product released during apoptosis) and circulating tumor DNA (ctDNA), offer complementary readouts of pathway engagement and early response. ctDNA-based monitoring in NSCLC is an area of active development, with applications in minimal residual disease detection and treatment response assessment . For rural settings, liquid biopsy approaches are particularly attractive because they reduce dependence on tissue biopsy infrastructure and can be performed at district-level laboratories with appropriate sample-handling protocols.

A trial design in which patients are stratified by functional apoptotic dependence—rather than by histology or a single mutation—is technically feasible and clinically necessary. Baseline expression of the Bcl-2 family member targeted by a given BH3 mimetic is a reasonable minimal enrollment criterion, provided it is interpreted alongside functional profiling rather than as a stand-alone predictor.

9. Implications for Rural Healthcare Advancement in India

The mechanistic literature reviewed above has a direct bearing on how apoptosis-targeted therapy should be deployed in rural India—and on why it currently is not.

9.1 The Rural Lung Cancer Burden

Rural India combines a high prevalence of smoking and smokeless tobacco use, widespread biomass-fuel exposure, and limited access to screening and molecular diagnostics, producing later-stage presentation and poorer outcomes than in

metropolitan centers (Indian Council of Medical Research, 2020; Gupta et al., 2011). The burden is not static: GBD 2021 data show that age-standardized lung cancer incidence and mortality rates in India have risen steadily, with the highest annual percentage changes in states including Uttarakhand, Uttar Pradesh, and Gujarat, many of which have large rural populations . At the same time, specialized thoracic oncology, immunohistochemistry, and next-generation sequencing are concentrated in urban tertiary centers, so that the molecular characterization on which apoptosis-targeted therapy depends is least available where the burden is greatest.

9.2 Four Linked Priorities for Rural-Relevant Translation

First, upstream risk reduction. Tobacco cessation, smokeless-tobacco control, and cleaner cooking-fuel transitions address the dominant preventable causes of rural lung cancer and reduce incidence at the population level (Gupta et al., 2011). These are the most cost-effective interventions available and should not be deferred in favor of downstream therapeutics.

Second, decentralized diagnostic pathways. Earlier detection and molecular characterization require decentralized sputum cytology, low-dose CT where feasible, telepathology for immunohistochemistry interpretation, and affordable biomarker testing. Low-dose CT screening has been shown to be feasible in India, with a lung cancer detection rate of 2.2% in a high-risk cohort and stage I diagnoses identified in three of four detected cases . Telepathology networks connecting district hospitals to regional cancer centers are a practical means of extending pathology expertise to rural settings, though standardization, infrastructure investment, and regulatory frameworks for AI-assisted image analysis remain works in progress . Without these, apoptosis-targeted therapy cannot be matched to the tumors that would benefit, and rural patients will continue to present at stages where curative-intent treatment is no longer possible.

Third, context-appropriate therapeutic delivery. Apoptosis-targeted agents must be developed and priced for delivery closer to home: oral BH3 mimetics and oral p53-reactivating compounds, regimens that can be safely administered in district day-care units, and tumor-selective delivery platforms that reduce toxicity and supportive-care burden. The cardiac toxicity observed with systemic MCL-1 inhibition is a particular concern in this context, because it requires monitoring infrastructure that may not be available at district level. This argues for prioritizing tumor-selective delivery platforms and oral agents with wider safety margins. The concentration of CAR-T and other advanced cellular therapies in apex centers is not a model that can be replicated at scale in rural India; the relevant question is which apoptosis-targeted approaches can be delivered safely outside tertiary care.

Fourth, biomarker validation in resource-limited settings. Functional biomarkers such as BH3 profiling and circulating apoptotic markers must be validated in rural and resource-limited populations, so that patient selection is based on tumor biology rather than on ability to pay or geographic access. DBP is technically demanding but can be performed on small tumor samples and has been adapted for high-throughput formats, suggesting that with appropriate quality-control frameworks it could be deployed at regional reference laboratories serving rural

catchment areas . Without such validation, precision oncology will widen rather than narrow the rural–urban survival gap.

9.3 Health-System Integration

These priorities are best implemented through a hub-and-spoke framework in which regional tertiary cancer centers provide molecular interpretation, protocol selection, and tele-oncology support, while district hospitals deliver oral targeted therapy, low-intensity combination regimens, and supportive care. Telepathology networks are a practical example of this architecture: digital slide scanning at district level, interpretation at regional hubs, and AI-assisted prescreening to prioritize cases, with the reporting pathologist retaining ultimate diagnostic responsibility . Community health workers can support tobacco cessation, early symptom recognition, and adherence monitoring. This architecture mirrors the decentralized models being developed for cancer care in India, but it has not been applied to apoptosis-targeted lung cancer therapy, and doing so is the central translational challenge for the coming decade.

10. Conclusion

Apoptosis dysregulation in lung cancer is a multi-layered problem: Bcl-2 family imbalance (with MCL-1 predominating in NSCLC), p53 inactivation, death-receptor pathway defects, and survival-signaling rewiring converge to produce a phenotype in which malignant cells survive therapeutic insults that would kill normal cells. The mechanistic literature has mapped these layers in detail, and the clinical literature has provided a complementary education in what happens when they are targeted in patients. Single-agent BH3 mimetics have limited activity in lung cancer; p53 reactivation has not yet produced durable benefit as monotherapy; and death-receptor agonists have been constrained by resistance and toxicity. Yet the same clinical experience has yielded design principles that are beginning to produce results: MCL-1-selective inhibition matched to NSCLC dependency, tumor-directed delivery of Bcl-2-family inhibitors, allele-specific p53 reactivation, and combinations that engage both apoptotic pathways and immune mechanisms.

For rural India, these principles carry particular urgency. Apoptosis-targeted therapies will not improve rural outcomes if they remain confined to urban tertiary centers. Rural-relevant translation requires linked progress in upstream risk reduction, decentralized diagnosis and molecular characterization, low-cost tumor-selective delivery, and validation of functional biomarkers in resource-limited settings. The cardiac toxicity of systemic MCL-1 inhibition and the technical demands of functional profiling are not arguments against pursuing apoptosis-targeted therapy in rural populations—they are arguments for designing delivery platforms and biomarker workflows that are compatible with rural health-system constraints. Understanding the molecular mechanisms behind apoptotic dysregulation remains essential, but the field has matured to the point where the more urgent task is to deploy those mechanisms selectively—in the right patients, with the right combination partners, at the right point in the therapeutic sequence, and in the right health-system context.

References

Acehan, D., Jiang, X., Morgan, D. G., Heuser, J. E., Wang, X., & Akey, C. W. (2002). Three-dimensional

structure of the apoptosome: Implications for assembly, procaspase-9 binding, and activation. *Molecular Cell*, 9(2), 423–432.

Ashkenazi, A. (2002). Targeting death and decoy receptors of the tumour-necrosis factor superfamily. *Nature Reviews Cancer*, 2(6), 420–430.

Ashkenazi, A. (2008). Directing cancer cells to self-destruct with pro-apoptotic receptor agonists. *Nature Reviews Drug Discovery*, 7(12), 1001–1012.

Ashkenazi, A., & Dixit, V. M. (1998). Death receptors: Signaling and modulation. *Science*, 281(5381), 1305–1308.

Aubrey, B. J., Strasser, A., & Kelly, G. L. (2016). Tumor-suppressor functions of the TP53 pathway. *Cold Spring Harbor Perspectives in Medicine*, 6(5), a026062.

Bratton, S. B., & Salvesen, G. S. (2010). Regulation of the Apaf-1-caspase-9 apoptosome. *Journal of Cell Science*, 123(19), 3209–3214.

Chen, L., Willis, S. N., Wei, A., Smith, B. J., Fletcher, J. I., Hinds, M. G., & Huang, D. C. (2005). Differential targeting of prosurvival Bcl-2 proteins by their BH3-only ligands allows complementary apoptotic function. *Cell Death and Differentiation*, 12(1), 112–125.

Cory, S., & Adams, J. M. (2002). The Bcl2 family: Regulators of the cellular life-or-death switch. *Nature Reviews Cancer*, 2(9), 647–656.

Deng, J., Carlson, N., Takeyama, K., Dal Cin, P., Shipp, M., & Letai, A. (2007). BH3 profiling identifies three distinct classes of apoptotic blocks to predict response to ABT-737 and conventional chemotherapeutic agents. *Cancer Cell*, 12(2), 171–185.

Dimberg, L. Y., Anderson, C. K., Camidge, R., Behbakht, K., Thorburn, A., & Ford, H. L. (2013). On the TRAIL to successful cancer therapy? Predicting and counteracting resistance against TRAIL-based therapeutics. *Oncogene*, 32(11), 1341–1350.

Djerbi, M., Screpanti, V., Catrina, A. I., Bogen, B., Biberfeld, P., & Grandien, A. (1999). The inhibitor of death receptor signaling, FLICE-inhibitory protein defines a new class of tumor progression factors. *The Journal of Experimental Medicine*, 190(7), 1025–1032.

Elmore, S. (2007). Apoptosis: A review of programmed cell death. *Toxicologic Pathology*, 35(4), 495–516.

Gonzalvez, F., & Ashkenazi, A. (2010). New insights into apoptosis signaling by Apo2L/TRAIL. *Oncogene*, 29(34), 4752–4765.

Green, D. R., & Kroemer, G. (2004). The pathophysiology of mitochondrial cell death. *Science*, 305(5684), 626–629.

Gross, A., McDonnell, J. M., & Korsmeyer, S. J. (1999). BCL-2 family members and the mitochondria in apoptosis. *Genes & Development*, 13(15), 1899–1911.

Gupta, P. C., Ray, C. S., Sinha, D. N., & Singh, P. K. (2011). Smokeless tobacco: A major public health problem in the SEA region: A

review. *Indian Journal of Public Health*, 55(3), 199–209.

Hanahan, D., & Weinberg, R. A. (2011). Hallmarks of cancer: The next generation. *Cell*, 144(5), 646–674.

Herbst, R. S., Eckhardt, S. G., Kurzrock, R., Ebbinghaus, S., O'Dwyer, P. J., Gordon, M. S., ... & Ashkenazi, A. (2010). Phase I dose-escalation study of recombinant human Apo2L/TRAIL, a dual proapoptotic receptor agonist, in patients with advanced cancer. *Journal of Clinical Oncology*, 28(17), 2839–2846.

Indian Council of Medical Research. (2020). *National Cancer Registry Programme Report 2020*. ICMR.

Kale, J., Osterlund, E. J., & Andrews, D. W. (2018). BCL-2 family proteins: Changing partners in the dance towards death. *Cell Death & Differentiation*, 25(1), 65–80.

Kischkel, F. C., Hellbardt, S., Behrmann, I., Germer, M., Pawlita, M., Krammer, P. H., & Peter, M. E. (1995). Cytotoxicity-dependent APO-1 (Fas/CD95)-associated proteins form a death-inducing signaling complex (DISC) with the receptor. *The EMBO Journal*, 14(22), 5579–5588.

Letai, A. (2008). Diagnosing and exploiting cancer's addiction to blocks in the apoptotic pathway. *Nature Reviews Cancer*, 8(2), 121–132.

Li, H., Zhu, H., Xu, C. J., & Yuan, J. (1998). Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis. *Cell*, 94(4), 491–501.

Luo, X., Budihardjo, I., Zou, H., Slaughter, C., & Wang, X. (1998). Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors. *Cell*, 94(4), 481–490.

Micheau, O., & Tschopp, J. (2003). Induction of TNF receptor I-mediated apoptosis via two sequential signaling complexes. *Cell*, 114(2), 181–190.

Moldoveanu, T., Grace, C. R., Llambi, F., Nourse, A., Fitzgerald, P., Gehring, K., & Green, D. R. (2013). BID-induced structural changes in BAK promote apoptosis. *Nature Structural & Molecular Biology*, 20(5), 589–597.

Peter, M. E., & Krammer, P. H. (2003). The CD95 (APO-1/Fas) DISC and beyond. *Cell Death & Differentiation*, 10(1), 26–35.

Ren, D., Tu, H. C., Kim, H., Wang, G. X., Bean, G. R., Takeuchi, O., ... & Cheng, E. H. (2010). BID, BIM, and PUMA are essential for activation of the BAX- and BAK-dependent cell death program. *Science*, 330(6009), 1390–1393.

Riedl, S. J., & Salvesen, G. S. (2007). The apoptosome: Signalling platform of cell death. *Nature Reviews Molecular Cell Biology*, 8(5), 405–413.

Roberts, A. W., Davids, M. S., Pagel, J. M., Kahl, B. S., Puvvada, S. D., Gerecitano, J. F., ... & Seymour, J. F. (2016). Targeting BCL2 with venetoclax in relapsed chronic lymphocytic leukemia. *New England Journal of Medicine*, 374(4), 311–322.

Roth, J. A., Nguyen, D., Lawrence, D. D., Kemp, B. L., Carrasco, C. H., Ferson, D. Z., ... & Cristiano, R. J. (1996). Retrovirus-mediated wild-type p53 gene transfer to tumors of patients with lung cancer. *Nature Medicine*, 2(9), 985–991.

Safa, A. R. (2012). c-FLIP, a master anti-apoptotic regulator. *Experimental Oncology*, 34(3), 176–184.

Souers, A. J., Leverson, J. D., Boghaert, E. R., Ackler, S. L., Catron, N. D., Chen, J., ... & Elmore, S. W. (2013). ABT-199, a potent and selective BCL-2 inhibitor, achieves antitumor activity while sparing platelets. *Nature Medicine*, 19(2), 202–208.

Vivanco, I., & Sawyers, C. L. (2002). The phosphatidylinositol 3-kinase AKT pathway in human cancer. *Nature Reviews Cancer*, 2(7), 489–501.

Vousden, K. H., & Lane, D. P. (2007). p53 in health and disease. *Nature Reviews Molecular Cell Biology*, 8(4), 275–283.

Vousden, K. H., & Prives, C. (2009). Blinded by the light: The growing complexity of p53. *Cell*, 137(3), 413–431.

Willis, S. N., Chen, L., Dewson, G., Wei, A., Naik, E., Fletcher, J. I., ... & Huang, D. C. (2005). Proapoptotic Bak is sequestered by Mcl-1 and Bcl-xL, but not Bcl-2, until displaced by BH3-only proteins. *Genes & Development*, 19(11), 1294–1305.

Youle, R. J., & Strasser, A. (2008). The BCL-2 protein family: Opposing activities that mediate cell death. *Nature Reviews Molecular Cell Biology*, 9(1), 47–59.

Journal of Rural Advancement
[ISSN 2347-2561 (P); 2583-6102 (E)]
Vol. 12, Issue 02, October 2024

Harnessing Plant–Microbe Interactions for Sustainable Agricultural Transformation in Rural India

Prateek Jain

Department of Biotechnology and Bioinformatics, North-Eastern Hill University, Shillong 793022, Meghalaya, India
Email: prateekjain20000@gmail.com
Received on: June 10, 2024 ; Accepted on: July 3, 2024

Abstract

Indian agriculture stands at a critical ecological and socioeconomic crossroads. While the Green Revolution enabled national food self-sufficiency, decades of intensive chemical fertilizer and pesticide application, monocropping, and groundwater depletion have precipitated severe soil organic carbon (SOC) loss, secondary salinization, declining factor productivity, and environmental degradation across rural agroecosystems. Concurrently, climate change has intensified drought, heatwaves, and erratic monsoons, disproportionately affecting smallholder and marginal farmers who cultivate over 86% of India's operational landholdings. Harnessing beneficial plant–microbe interactions—encompassing plant growth-promoting rhizobacteria (PGPR), arbuscular mycorrhizal fungi (AMF), nitrogen-fixing diazotrophs, endophytes, and biological control agents—offers an ecologically sound, economically viable pathway to restore soil health, enhance nutrient-use efficiency (NUE), confer abiotic and biotic stress resilience, and reduce agrochemical dependency. This review synthesizes the mechanistic foundations of rhizosphere and endosphere symbioses, evaluates region-specific translational opportunities across India's diverse agro-climatic zones (from the Indo-Gangetic Plains and rainfed drylands to the acidic soils of the North-Eastern Himalayas), and critically examines the biological, technological, regulatory, and socioeconomic bottlenecks hindering large-scale rural adoption. Finally, we outline a forward-looking roadmap integrating multi-omics-guided synthetic microbial communities (SynComs), nano-encapsulated and biofilm-based bioformulations, farmer producer organization (FPO)-led decentralized biomanufacturing, and supportive policy reforms to drive a second, evergreen agricultural transformation in rural India.

Keywords: Plant–microbe interactions; Biofertilizers; Plant growth-promoting rhizobacteria (PGPR); Arbuscular mycorrhizal fungi (AMF); Sustainable agriculture; Smallholder farmers; Rural India; Synthetic microbial communities (SynComs)

1. Introduction

Agriculture remains the socioeconomic backbone of rural India, supporting the livelihoods of nearly half of the nation's workforce and contributing approximately 18% to the gross domestic product (GDP). Following the Green Revolution of the late 1960s, the widespread adoption of high-yielding crop varieties, assured irrigation, and synthetic agrochemicals propelled India from chronic food deficits to a global agricultural powerhouse (Pingali, 2012). However, the long-term ecological externalities of this input-intensive paradigm have become increasingly unsustainable. Decades of imbalanced

nitrogen–phosphorus–potassium (NPK) fertilization—heavily skewed toward subsidized urea—have depleted soil organic matter, disrupted native soil microbial biodiversity, acidified or salinized arable soils, and caused widespread multi-nutrient deficiencies, particularly in zinc, iron, boron, and sulfur (Bijay-Singh & Craswell, 2021; Bhattacharyya et al., 2015). Furthermore, the partial factor productivity of synthetic fertilizers in major cropping systems such as the rice–wheat rotation of the Indo-Gangetic Plains has declined precipitously, forcing farmers to apply escalating doses of chemical inputs merely to maintain baseline yields, thereby eroding farm profitability and contaminating groundwater with nitrates and pesticide residues (Ladha et al., 2003; Abrol et al., 2017).

These soil degradation challenges are compounded by the structural realities of rural Indian farming and accelerating climate change. Approximately 86.2% of Indian farmers are smallholders or marginal cultivators owning less than two hectares of land, operating with limited capital reserves and high vulnerability to input price volatility and climatic shocks (Birthal et al., 2015). Nearly 51% of India's net sown area remains rainfed, exposing crops to recurrent droughts, terminal heat stress, soil salinity, and emerging pest and pathogen outbreaks (Venkateswarlu & Shanker, 2009). Simultaneously, heavy reliance on imported potassic and phosphatic raw materials places a massive fiscal burden on the national exchequer through fertilizer subsidies, underscoring the urgent strategic imperative for domestic, bio-based alternatives (Sapkota & Bijay-Singh, 2025).

In nature, plants do not function as autonomous entities; rather, they operate as integrated ecological units—termed holobionts—co-evolving with complex communities of bacteria, fungi, archaea, and protists colonizing the rhizosphere (root–soil interface), rhizoplane (root surface), endosphere (internal tissues), and phyllosphere (aerial surfaces) (Vandenkoornhuyse et al., 2015; Bulgarelli et al., 2013). Through sophisticated chemical signaling mediated by root exudates—comprising organic acids, sugars, amino acids, flavonoids, and strigolactones—plants actively recruit beneficial microbiota that reciprocate by mobilizing locked soil nutrients, fixing atmospheric nitrogen, synthesizing phytohormones, priming systemic immunity, and buffering abiotic stresses (Bais et al., 2006; Sasse et al., 2018). For rural India, translating these plant–microbe interactions into low-cost, locally adapted biofertilizers, biostimulants, and biopesticides represents a transformative opportunity to decouple agricultural productivity from ecological degradation (Gupta et al., 2021; Singh et al., 2016).

Despite decades of foundational microbiological research across Indian agricultural universities and national institutes, the field-level translation of microbial technologies remains uneven. Commercial biofertilizers currently substitute only a fraction of chemical fertilizer demand, constrained by inconsistent field efficacy across heterogeneous agro-climatic zones, short shelf life under tropical storage conditions, suboptimal carrier formulations, quality control deficits, and weak last-mile extension networks (Vessey, 2003; Herrmann & Lesueur, 2013; Kumar et al., 2021). This review provides a comprehensive, critical synthesis of the mechanisms governing beneficial plant–microbe symbioses, evaluates their agronomic potential across India's major rural agroecosystems, dissects the technical and institutional barriers to adoption, and proposes next-generation biotechnological and policy interventions to mainstream

microbiome-driven sustainable agriculture in rural India.

2. Mechanistic Foundations of Beneficial Plant–Microbe Interactions

Beneficial plant-associated microorganisms—broadly categorized into plant growth-promoting rhizobacteria (PGPR), nitrogen-fixing diazotrophs, arbuscular mycorrhizal fungi (AMF), plant growth-promoting fungi (PGPF), and endophytes—facilitate plant growth and health via interconnected direct and indirect physiological pathways (Glick, 2012; Bhattacharyya & Jha, 2012) (Table 1).




Figure-1. Ectomycorrhizal vs. endomycorrhizal root colonization.

As illustrated in the mycorrhizal colonization diagram above, endomycorrhizal fungi penetrate root cortical cells to form branching arbuscules and external hyphae that expand nutrient absorption beyond the root zone, complementing rhizosphere bacteria and soil-borne antagonists across the soil–root continuum.

2.1 Nutrient Acquisition and Biogeochemical Cycling

Nitrogen (N) and phosphorus (P) are the primary limiting macronutrients across Indian agricultural soils, yet conventional synthetic applications suffer from dismal recovery efficiencies—typically 30–40% for applied N and only 15–20% for applied P (Tilman et al., 2002). Biological nitrogen fixation (BNF) represents the most sustainable alternative to synthetic Haber–Bosch nitrogen. Symbiotic rhizobia (*Rhizobium*, *Bradyrhizobium*, *Mesorhizobium*, *Sinorhizobium*) establish specialized root nodules in leguminous crops (chickpea, pigeonpea, mungbean, blackgram, groundnut, soybean), wherein host-secreted flavonoids activate bacterial NodD regulators, triggering Nod factor synthesis and root hair curling (Oldroyd, 2013; Herridge et al., 2008). Inside oxygen-buffered symbiosomes, bacteroids express the nitrogenase complex to reduce atmospheric N_2 into bioavailable ammonia. In non-leguminous cereals (rice, wheat, maize, millets) and cash crops like sugarcane, associative and endophytic diazotrophs—including *Azospirillum brasilense*, *Azotobacter chroococcum*, *Herbaspirillum seropedicae*, and *Gluconacetobacter diazotrophicus*—colonize root surfaces and intercellular spaces, contributing substantial fixed nitrogen while minimizing leaching and

nitrous oxide (N_2O) emissions (Boddey et al., 2003; Santi et al., 2013).

Functional Category	Key Microbial Taxa	Primary Biochemical Mechanisms	Agronomic Outcomes in Indian Crops	Representative References
Symbiotic & Associative N_2 Fixers	*Rhizobium*, *Bradyrhizobium*, *Mesorhizobium*, *Azospirillum*, *Azotobacter*, *Gluconacetobacter*	Reduction of atmospheric NH_2 to NH_3 via oxygen-sensitive nitrogenase enzyme complex (*nif* genes)	Supplies 20–40 kg N/ha in cereals/sugarcane and up to 100–150 kg N/ha in pulses/groundnut	Herridge et al. (2008); Boddey et al. (2003); Santi et al. (2013)
Phosphate & Potassium Solubilizers	*Bacillus megaterium*, *Pseudomonas striata*, *Pantoea*, *Aspergillus niger*, *Penicillium*, *Frateuria aurantia*	Secretion of gluconic, citric, and oxalic acids; release of acid/alkaline phosphatases and phytases	Unlocks insoluble Ca-, Fe-, and Al-bound phosphorus in alkaline and acidic Indian soils	Richardson & Simpson (2011); Alori et al. (2017); Meena et al. (2014)
Mycorrhizal Symbionts	*Funneliformis mosseae*, *Rhizophagus irregularis*, *Gigaspora*, *Glomus* spp.	Extraradical hyphal network expansion; glomalin-related soil protein (GRSP) secretion; polyphosphate transport	Enhances P, Zn, and water uptake by 30–60%; stabilizes soil aggregates in drylands	Smith & Read (2008); Rillig (2004); Auge (2001)
Phytohormone & ACC Deaminase Producers	*Pseudomonas fluorescens*, *Bacillus subtilis*, *Enterobacter*, *Azospirillum brasilense*	Biosynthesis of IAA, gibberellins, and cytokinins; cleavage of ACC into α-ketobutyrate and NH_3	Stimulates lateral root proliferation; lowers stress-induced ethylene under drought and salinity	Glick et al. (2007); Spaepen et al. (2007)
Biological Control Agents (BCAs)	*Trichoderma harzianum*, *T. viride*, *Pseudomonas fluorescens*, *Bacillus amyloliquefaciens*, *Streptomyces*	Mycoparasitism (chitinases, β-1,3-glucanases), antibiosis (2,4-DAPG, phenazines, lipopeptides), ISR priming	Suppresses *Fusarium* wilt, *Rhizoctonia* root rot, *Pythium* damping-off, and bacterial blight	Harman et al. (2004); Haas & Défago (2005); Pieterse et al. (2014)

Phosphorus in Indian soils is predominantly immobilized as insoluble tricalcium phosphate in alkaline/calcareous soils (Indo-Gangetic Plains and Deccan Plateau) or bound to iron and aluminum oxides in acidic soils (North-Eastern and Eastern India). Phosphate-solubilizing microorganisms (PSMs)—including *Bacillus megaterium*, *Pseudomonas putida*, *Enterobacter*, Aspergillus, and Penicillium—solubilize inorganic phosphates by secreting low-molecular-weight organic acids (gluconic, 2-ketogluconic, citric, malic, and oxalic

acids) that chelate Ca2+, Fe3+, and Al3+ cations and lower rhizosphere pH (Rodríguez & Fraga, 1999; Alori et al., 2017). Concurrently, they mineralize organic phosphorus pools via extracellular phytases and acid/alkaline phosphor-monoesterases (Richardson & Simpson, 2011). Similarly, potassium-solubilizing bacteria (Frateuria aurantia, Bacillus mucilaginosus) weather silicate minerals such as mica and feldspar via acidolysis and capsular polysaccharide complexation, reducing India's dependence on imported muriate of potash (Meena et al., 2014).

Arbuscular mycorrhizal fungi (phylum Glomeromycota, e.g., *Rhizophagus irregularis*, *Funneliformis mosseae*) form obligate endosymbioses with over 80% of terrestrial crop species. Upon perceiving host-derived strigolactones, AMF hyphae penetrate root cortical cells to form highly branched arbuscules—specialized interfaces for bidirectional nutrient exchange—while extending a vast extraradical hyphal network far beyond the root phosphorus-depletion zone (Smith & Read, 2008; Parniske, 2008). Beyond phosphorus, AMF hyphae efficiently scavenge immobile micronutrients such as zinc (Zn) and copper (Cu), directly addressing widespread micronutrient malnutrition ("hidden hunger") in Indian staple grains (Lehmann et al., 2014; Singh & Prasanna, 2020). Furthermore, high-affinity microbial siderophores (catecholates, hydroxamates, and carboxylates) produced by *Pseudomonas* and *Bacillus* chelate insoluble ferric iron (Fe^{3+}) in calcareous soils, facilitating plant iron acquisition while starving phytopathogens of essential iron (Kloepper et al., 1980; Saha et al., 2016).

2.2 Phytohormonal Modulation and Root System Architecture

Beneficial microbes profoundly remodel plant root architecture—increasing lateral root density, root hair elongation, and total absorptive surface area—by modulating endogenous phytohormone homeostasis (Spaepen et al., 2007). Over 80% of rhizosphere bacteria synthesize indole-3-acetic acid (IAA) primarily via the tryptophan-dependent indole-3-pyruvate (IPyA) pathway; low-to-moderate bacterial IAA concentrations stimulate primary root elongation and lateral root primordia initiation, enhancing water and nutrient foraging in nutrient-poor soils (Spaepen et al., 2007; Patten & Glick, 2002). Additionally, PGPR and PGPF produce gibberellins (promoting stem elongation and seed germination) and cytokinins (stimulating cell division and delaying leaf senescence) (Arkhipova et al., 2005).

A pivotal mechanism under environmental stress is the enzymatic cleavage of 1-aminocyclopropane-1-carboxylate (ACC), the immediate biosynthetic precursor of ethylene in higher plants. Under drought, salinity, waterlogging, or pathogen attack, plants synthesize deleterious bursts of "stress ethylene" that inhibit root elongation and accelerate senescence. PGPR possessing the enzyme **ACC deaminase** (encoded by *acdS*, common in *Pseudomonas*, *Burkholderia*, *Variovorax*, and *Enterobacter*) sequester exuded ACC from plant roots and hydrolyze it into ammonia and α-ketobutyrate, thereby dampening ethylene accumulation and sustaining root growth under adverse field conditions (Glick et al., 2007; Glick, 2014).

2.3 Abiotic Stress Mitigation: Drought, Salinity, and Temperature Extremes

Climate-induced abiotic stresses severely constrain crop yields across rural India,

where ~6.73 million hectares are affected by salinity and sodicity and over two-thirds of agricultural land is prone to periodic drought (Sharma et al., 2016). Halotolerant and drought-adapted PGPR and AMF confer systemic abiotic stress tolerance through coordinated physiological interventions (Yang et al., 2009; Dodd & Pérez-Alfocea, 2012). First, microbes trigger osmotic adjustment by inducing plant accumulation of compatible solutes—proline, glycine betaine, trehalose, and soluble sugars—that stabilize proteins and cellular membranes without disrupting enzymatic activity (Bano & Fatima, 2009; Kaushal & Wani, 2016). Second, under saline conditions, beneficial rhizobacteria regulate host ion transporters (such as the high-affinity potassium transporter *HKT1* and salt overly sensitive *SOS1* antiporters), restricting toxic Na^+ influx into shoots while maintaining a favorable cytosolic $K^+\backslash Na^+$ ratio (Zhang et al., 2008; Bharti et al., 2016).

Third, microbial inoculation upregulates plant antioxidant defense enzymes—including superoxide dismutase (SOD), catalase (CAT), ascorbate peroxidase (APX), and glutathione reductase (GR)—scavenging cytotoxic reactive oxygen species (ROS) generated during drought and heat stress (Gururani et al., 2013). Fourth, bacterial exopolysaccharides (EPS) form hydrophilic biofilms around root surfaces, binding Na^+ ions, enhancing rhizosheath water retention, and improving soil macro-aggregation alongside AMF-secreted glomalin-related soil proteins (GRSP) (Sandhya et al., 2009; Rillig, 2004; Auge, 2001).

2.4 Biocontrol of Phytopathogens and Induced Systemic Resistance

Soil-borne fungal, bacterial, and nematode pathogens—including *Fusarium oxysporum*, *Rhizoctonia solani*, *Macrophomina phaseolina*, *Sclerotium rolfsii*, *Xanthomonas oryzae*, and root-knot nematodes (*Meloidogyne* spp.)—inflict annual crop losses of 15–25% in India. Biological control agents (BCAs) suppress disease through both direct antagonism and host immune priming (Compant et al., 2005; Köhl et al., 2019). *Trichoderma* species (*T. harzianum*, *T. viride*, *T. asperellum*), which account for the majority of commercial biofungicides in India, exhibit aggressive mycoparasitism: they sense pathogen-derived oligochitins, coil around host hyphae, form appressoria, and secrete cell wall-degrading enzymes (chitinases, β-1,3-glucanases, and proteases) alongside antifungal peptaibols, gliotoxin, and plant-derived bioactive compounds with complementary antibacterial and antifungal properties (Harman et al., 2004; Druzhinina et al., 2011; Tyśkiewicz et al., 2022; Sharma, 2024).

Fluorescent pseudomonads (*P. fluorescens*, *P. protegens*) and *Bacillus* species (*B. subtilis*, *B. amyloliquefaciens*, *B. velezensis*) produce broad-spectrum antimicrobial metabolites, including 2,4-diacetylphloroglucinol (2,4-DAPG), phenazines, pyrrolnitrin, pyoluteorin, hydrogen cyanide (HCN), and cyclic lipopeptides (surfactin, iturin, fengycin) that disrupt pathogen membrane integrity (Haas & Défago, 2005; Ongena & Jacques, 2008). Crucially, colonization by non-pathogenic PGPR and *Trichoderma* triggers Induced Systemic Resistance (ISR) throughout the entire plant. Unlike pathogen-induced systemic acquired resistance (SAR), which relies on salicylic acid (SA) and pathogenesis-related (PR) proteins, ISR is predominantly mediated by jasmonic acid (JA) and ethylene (ET) signaling cascades, priming the host for

rapid callose deposition, phenolic accumulation, and phytoalexin synthesis upon subsequent attack by foliar fungi, bacteria, or insect herbivores (Pieterse et al., 2014; Van Wees et al., 2008).

3. Opportunities Across India's Diverse Rural Agroecosystems

India's vast pedoclimatic heterogeneity—spanning 15 distinct agro-climatic zones—necessitates tailored microbial interventions aligned with regional cropping systems, soil chemistry, and farmer socioeconomic profiles (Table 2).

3.1 Revitalizing the Indo-Gangetic Plains and Curbing Stubble Burning

The Indo-Gangetic Plains (IGP), often termed India's breadbasket, suffer from severe soil fatigue induced by continuous rice–wheat monocropping and excessive chemical fertilizer application (Ladha et al., 2003; Bhattacharyya et al., 2015). Integrated Nutrient Management (INM) combining 75% recommended NPK with composite biofertilizers (*Azotobacter chroococcum*, phosphate-solubilizing *Bacillus megaterium*, *Rhizobium*, and potassium-solubilizing *Frateuria aurantia*) consistently sustains or increases grain yields while restoring soil microbial biomass carbon and dehydrogenase activity (Mader et al., 2011; Kumar et al., 2021; Kumar et al., 2024).

Agro-Climatic Region	**Dominant Cropping Systems**	**Key Soil & Climatic Constraints**	**Target Microbial Interventions**	**Documented Agronomic & Ecological Benefits**
Indo-Gangetic Plains (IGP)	Rice–wheat, rice–maize, sugarcane	Declining SOC, alkaline pH (7.8–8.5), P fixation, groundwater depletion, crop residue burning	*Azotobacter* + PSB + KSB consortia; *Trichoderma* & lignocellulolytic fungi (*Aspergillus*, *Phanerochaete*) for in situ straw decomposition	15–25% reduction in synthetic NPK; rapid 20–25 day paddy straw biodecomposition, curbing stubble burning
Rainfed & Semi-Arid Deccan / Central India	Pulses (chickpea, pigeonpea), oilseeds (groundnut, soybean), cotton, millets	Moisture deficit, terminal drought, low soil organic matter, *Fusarium* wilt, dry root rot	Native drought-tolerant *Rhizobium* + AMF (*F. mosseae*) + ACC-deaminase *Pseudomonas* + *T. harzianum*	12–28% yield gain under rainfed conditions; enhanced nodulation, P/Zn uptake, and wilt suppression
North-Eastern & Himalayan Hill Regions	Upland rice, maize, ginger, turmeric, citrus, tea, vegetables	Strong soil acidity (pH 4.5–5.5), Al/Fe toxicity, P immobilization, steep slope erosion	Acid-tolerant PSB (*Burkholderia*, *Penicillium*), AMF, *Azospirillum*, and indigenous *Trichoderma* strains	Mitigates Al toxicity; unlocks Fe/Al-bound P; supports certified organic farming (e.g., Sikkim model)
Coastal & Arid Salt-Affected Belts	Salt-tolerant rice, cotton, mustard, pearl millet, horticulture	Primary/secondary soil salinity and sodicity (ECe greater than 4 dS/m), high Na^+ toxicity	Halotolerant PGPR (*Bacillus licheniformis*, *Halomonas*, *Pseudomonas*) + salt-adapted AMF	Improves K^+/Na^+ selectivity, osmolyte accumulation, and grain yield by 15–30% on marginal saline lands

Beyond nutrient supply, microbial consortia offer an immediate solution to the acute environmental crisis of paddy straw burning in Punjab, Haryana, and western Uttar Pradesh. Application of lignocellulolytic microbial decomposers—such as the Pusa Decomposer consortium comprising *Aspergillus awamori*, *Trichoderma viride*, *Phanerochaete chrysosporium*, and *Aspergillus nidulans*—accelerates in situ breakdown of silica-rich rice residues within 20–25 days, recycling ~6–7 kg N, 1–1.5 kg P, and 15–20 kg K per tonne of straw back into the soil while eliminating particulate air pollution (Kumar et al., 2008; Chaudhary et al., 2017).

3.2 Climate-Proofing Rainfed Drylands and Pulse-Oilseed Systems

Over 80% of India's pulses, oilseeds, and coarse cereals (sorghum, pearl millet, finger millet) are cultivated in rainfed drylands characterized by erratic rainfall and impoverished soils (Venkateswarlu & Shanker, 2009). Dual inoculation of legumes such as chickpea (*Cicer arietinum*), pigeonpea (*Cajanus cajan*), and green gram (*Vigna radiata*) with host-specific *Mesorhizobium*/*Bradyrhizobium*/*Rhizobium*, phosphate-solubilizing bacteria, and arbuscular mycorrhizal fungi (*Funneliformis mosseae*) generates strong tripartite synergy: AMF improve phosphorus and water supply required for energy-intensive nitrogenase activity, while rhizobia and PSB enhance root colonization, phosphorus bioavailability, and nitrogen nutrition, boosting grain yields by 15–30% even under moisture-limited regimes (Mortimer et al., 2008; Teja et al., 2022; Calderon & Dangi, 2024; Mugabo et al., 2024). Furthermore, seed biopriming of rainfed crops with ACC-deaminase-producing *Pseudomonas* and *Trichoderma harzianum* enhances seedling vigor, deep rooting, and tolerance against dry root rot (*Macrophomina phaseolina*) (Glick et al., 2007; Harman et al., 2004).

3.3 Advancing Organic and Natural Farming in the Himalayas and Tribal Belts

North-Eastern India and Himalayan hill states possess inherently high soil organic matter and low baseline chemical fertilizer usage, positioning them as natural hubs for certified organic agriculture—exemplified by Sikkim's transition to 100% organic farming (Das et al., 2017). However, strong soil acidity (pH 4.5–5.5) in these regions causes severe phosphorus fixation by iron and aluminum oxides and aluminum phytotoxicity. Bioprospecting indigenous acid-tolerant phosphate-solubilizing bacteria (*Burkholderia*, *Pantoea*), acid-adapted AMF, and native *Trichoderma* isolates from Himalayan rhizospheres unlocks fixed phosphorus and protects high-value spice and horticultural crops (ginger, turmeric, large cardamom, mandarin) from devastating rhizome rot and wilt diseases without synthetic agrochemicals (Pandey et al., 2006; Negi et al., 2025).

Similarly, microbial science provides an empirical foundation to validate and optimize traditional Indian agroecological practices—such as *Jeevamrutha* and *Beejamrutha* promoted under Zero Budget Natural Farming (ZBNF) / Bharatiya Prakritik Krishi Paddhati (BPKP). While traditional fermented formulations introduce diverse native *Bacillus*, *Pseudomonas*, lactic acid bacteria, and yeasts that stimulate soil enzymatic activity, scientific standardization and enrichment with verified elite PGPR/BCA strains can transform these low-cost on-farm preparations into reliable, high-potency bio-inputs for resource-poor tribal and marginal farmers (Smith et al., 2020; Negi et al., 2025).

3.4 Biofortification and Phytoremediation of Degraded Lands

Micronutrient deficiencies—particularly zinc and iron—affect both crop productivity across ~36–49% of Indian soils and human nutrition in rural populations (Alloway, 2009). Zinc-solubilizing bacteria (*Bacillus aryabhattai*, *Pseudomonas* spp.) and AMF mobilize insoluble zinc compounds (ZnO, $ZnCO_3$) via organic acid secretion and enhance root-to-grain translocation, achieving agronomic biofortification of wheat, rice, and maize grains alongside higher yields (Ramesh et al., 2014; Singh & Prasanna, 2020). Additionally, heavy-metal-tolerant PGPR and mycorrhizae immobilize or detoxify arsenic, cadmium, and chromium in contaminated peri-urban and alluvial soils (such as arsenic-affected groundwater belts of West Bengal and Bihar) through biosorption, siderophore chelation, and altered redox speciation, safeguarding rural food chains (Saha et al., 2016; Glick, 2010).

4. Key Challenges Hindering Field-Scale Translation in Rural India

Despite overwhelming laboratory and greenhouse evidence, the transition of plant–microbe technologies from research stations to millions of smallholder fields in rural India faces multi-dimensional bottlenecks spanning biology, manufacturing, regulation, and extension (Herrmann & Lesueur, 2013; Bashan et al., 2014; Kumar et al., 2021).

4.1 Ecological Inconsistency and the "Lab-to-Field" Performance Gap

A major scientific barrier is the poor reproducibility of single-strain microbial inoculants across diverse field environments. Strains selected under sterile, nutrient-controlled laboratory conditions frequently fail to establish in complex native soils due to intense competition with indigenous microflora, predation by bacterivorous protozoa and bacteriophages, incompatible soil pH, high ambient temperatures, or mismatched host crop genotypes (Bulgarelli et al., 2013; Sessitsch et al., 2019). Moreover, excessive chemical fertilizer application in intensively farmed regions actively suppresses symbiotic signaling: high soil nitrate inhibits legume nodulation, while elevated soluble phosphate downregulates strigolactone exudation and reduces plant carbon allocation to AMF, rendering inoculants agronomically redundant when applied alongside full chemical doses (Parniske, 2008; Calderon & Dangi, 2024).

4.2 Formulation, Shelf Life, and Tropical Supply Chain Constraints

Unlike synthetic chemicals, biofertilizers and biopesticides contain living cells or propagules whose viability declines rapidly under rural Indian storage and transport conditions, where ambient temperatures routinely exceed 35–45 °C without cold-chain infrastructure (Bashan et al., 2014). Historically, Indian biofertilizer production relied heavily on solid lignite, charcoal, or talc carriers, which suffer from high contamination rates, batch-to-batch physicochemical variability, short shelf life (often less than 3–6 months), and difficulty in mechanized seed or drip application (Herrmann & Lesueur, 2013; Malusá et al., 2012). While liquid biofertilizers formulated with osmoprotectants (trehalose, glycerol, polyvinylpyrrolidone) have improved viability to 12–18 months, maintaining high colony-forming unit (CFU) counts ($10^8\text{–}10^9\text{ CFU/mL}$) during last-mile village retail distribution remains a persistent logistical hurdle (Kumar et al., 2021; Negi et al., 2025).

4.3 Quality Control Deficits and Regulatory Fragmentation

Farmer trust is the single most fragile asset in rural input markets. Substandard commercial products—containing low viable cell counts, dead inocula, or pathogenic contaminants—have periodically flooded local markets, causing field failures that disillusion farmers and trigger reversion to chemical inputs (Herrmann & Lesueur, 2013; Vaidyanathan, 2022). Although biofertilizers are regulated under India's Fertilizer (Inorganic, Organic or Mixed) Control Order (FCO), 1985, and biopesticides under the Insecticides Act, 1968 (Central Insecticides Board and Registration Committee, CIBRC), enforcement capacity at the district level is severely constrained. State soil and biofertilizer testing laboratories often lack molecular identification facilities (such as 16S rRNA/ITS sequencing or qPCR) to distinguish active functional strains from inert contaminants, relying instead on slow, culture-dependent plate counts (Vaidyanathan, 2022). Furthermore, multi-strain microbial consortia and dual-purpose products (exhibiting both biofertilizer and biopesticidal traits) encounter overlapping, protracted regulatory approval pathways that discourage private and startup innovation (Negi et al., 2025).

4.4 Socioeconomic, Behavioral, and Extension Barriers

Smallholder farmers in rural India exhibit strong risk aversion, as a single crop failure can precipitate severe household debt. Chemical fertilizers deliver rapid, visible vegetative greening within days, whereas microbial inoculants act gradually through root colonization and soil biological activation, yielding subtler early visual cues (Bashan et al., 2014; Kumar et al., 2021). Additionally, massive public subsidies on urea distort market price signals, making synthetic nitrogen artificially cheap compared to unsubsidized quality biostimulants (Sapkota & Bijay-Singh, 2025). Compounding this economic asymmetry is an acute last-mile knowledge gap: many farmers and local agri-input dealers lack training on proper seed inoculation techniques, compatibility with chemical seed treatments (e.g., fungicides that kill *Rhizobium* or *Trichoderma* upon direct contact), soil moisture requirements at sowing, and expiry verification (Herrmann & Lesueur, 2013).

5. Future Perspectives and Strategic Roadmap

Overcoming these constraints requires converging frontier microbiome science with frugal formulation engineering, decentralized rural biomanufacturing, and coherent policy reform.

5.1 Multi-Omics-Guided Synthetic Microbial Communities (SynComs)

Rather than relying on isolated single strains, the future of agricultural microbiome engineering lies in designing Synthetic Microbial Communities (SynComs)—rationally assembled, functionally complementary consortia combining nitrogen fixers, P/K solubilizers, AMF, and biocontrol agents that mimic natural core microbiomes (Vorholt et al., 2017; Sessitsch et al., 2019). Integrating high-throughput amplicon sequencing, shotgun metagenomics, metatranscriptomics, and metabolomics enables researchers to map indigenous rhizosphere networks across India's agro-climatic zones and identify keystone taxa adapted to local soil pH, moisture, and crop cultivars (Bulgarelli et al., 2013; Sahil et al., 2025). Machine learning and genome-scale

metabolic modeling can predict cross-feeding synergies and ecological stability prior to field deployment, ensuring that introduced SynComs coexist harmoniously with native soil microbiota and persist across the crop lifecycle (Vorholt et al., 2017; Gupta et al., 2021).

5.2 Host-Microbiome Co-Breeding ("Microbiome-Smart" Crops)

Modern crop breeding during the Green Revolution inadvertently selected for high yield under luxury synthetic fertilizer regimes, often eroding wild plant traits governing root exudation and beneficial microbial recruitment (Pérez-Jaramillo et al., 2016). Future Indian crop improvement programs—leveraging genome-wide association studies (GWAS) and marker-assisted selection—must incorporate "microbiome-recruitment traits" (such as optimal flavonoid and strigolactone exudation profiles and root hair architecture) into elite varieties of rice, wheat, pulses, and millets (Pérez-Jaramillo et al., 2016; Gupta et al., 2020). Breeding cultivars that actively nurture beneficial rhizosphere partners maximizes biological nutrient acquisition and stress resilience without requiring repeated high-dose inoculation.

5.3 Next-Generation Bioformulations: Biofilms, Nano-Encapsulation, and Seed Coating

To conquer tropical shelf-life and field survival barriers, formulation science must transition from crude carrier powders to advanced delivery matrices (Bashan et al., 2014; Malusá et al., 2012). Biofilm-based biofertilizers—wherein beneficial bacteria (e.g., *Azotobacter* or *Bacillus*) are co-cultured with fungal partners (*Trichoderma* or *Aspergillus*) to form robust, extracellular polymeric matrix-embedded communities—exhibit superior desiccation tolerance, UV resistance, and root colonization efficiency compared to planktonic cells (Seneviratne et al., 2008; Kumar et al., 2021). Similarly, **polymeric micro- and nano-encapsulation** using biodegradable biopolymers (sodium alginate, chitosan, starch, carboxymethyl cellulose) enriched with trehalose or humic acids shields microbes from high storage temperatures and agrochemical toxicity while enabling controlled, sustained release in the rhizosphere (Schoebitz et al., 2013; Kumar et al., 2021). Industrial and FPO-level **film-coating of seeds** with stable microbial consortia prior to sale eliminates on-farm application errors and guarantees uniform rhizosphere establishment from germination onward (O'Callaghan, 2016).

5.4 Decentralized Rural Biomanufacturing and Policy Alignment

Scaling microbial agriculture across 140 million Indian farm holdings cannot rely solely on centralized urban factories. Establishing decentralized, village- or block-level bio-input resource centers (BRCs) operated by Farmer Producer Organizations (FPOs), women's self-help groups (SHGs), and rural agri-entrepreneurs—supported by state agricultural universities (SAUs) and Krishi Vigyan Kendras (KVKs) providing certified mother cultures and quality testing—can dramatically reduce cold-chain costs, create rural green employment, and supply fresh, region-specific inoculants (Vaidyanathan, 2022; Sapkota & Bijay-Singh, 2025).

At the national policy level, integrating biofertilizer recommendations into India's Soil Health Card (SHC) scheme and digital public infrastructure (AgriStack) can shift farmers from blanket chemical applications to precision Integrated Nutrient

Management (Sapkota & Bijay-Singh, 2025). Reforming fertilizer subsidy architectures—such as introducing direct benefit transfers (DBT) or nutrient-based incentives that reward soil organic carbon accrual and verified biofertilizer adoption—will level the economic playing field. Simultaneously, mandating QR-code-based batch traceability, molecular quality certification, and harmonized single-window regulatory clearance for multi-functional microbial consortia will safeguard product efficacy and build enduring farmer confidence (Vaidyanathan, 2022; Negi et al., 2025).

6. Conclusion

Harnessing beneficial plant–microbe interactions represents a cornerstone strategy for reconciling agricultural productivity, environmental stewardship, and smallholder prosperity in rural India. Decades of mechanistic research confirm that PGPR, nitrogen-fixing diazotrophs, arbuscular mycorrhizal fungi, and biocontrol fungi can substantially reduce synthetic fertilizer and pesticide requirements while buffering crops against drought, salinity, and soil-borne diseases. Realizing this potential at scale, however, demands bridging the persistent gap between laboratory discovery and rural field reality. By uniting multi-omics-guided SynCom design and host-microbiome co-breeding with heat-stable polymeric and biofilm formulations, decentralized FPO-led biomanufacturing, rigorous molecular quality enforcement, and subsidy realignment, India can transition from chemically intensive farming to a resilient, regenerative, and self-reliant rural agricultural economy.

References

Abrol, Y. P., Adhya, T. K., Aneja, V. P., Raghuram, N., Pathak, H., Kulshrestha, U., Sharma, C., & Singh, B. (2017). *The Indian Nitrogen Assessment: Sources of Reactive Nitrogen, Environmental and Climate Effects, Management Options, and Policies*. Elsevier.

Alloway, B. J. (2009). Soil factors associated with zinc deficiency in crops and humans. *Environmental Geochemistry and Health*, 31(5), 537–548.

Alori, E. T., Glick, B. R., & Babalola, O. O. (2017). Microbial phosphorus solubilization and its potential for use in sustainable agriculture. *Frontiers in Microbiology*, 8, 971.

Arkhipova, T. N., Veselov, S. U., Melentiev, A. I., Martynenko, E. V., & Kudoyarova, G. R. (2005). Ability of bacterium *Bacillus subtilis* to produce cytokinins and to influence the growth and endogenous hormone content of lettuce plants. *Plant and Soil*, 272(1), 201–209.

Auge, R. M. (2001). Water relations, drought and vesicular-arbuscular mycorrhizal symbiosis. *Mycorrhiza*, 11(1), 3–42.

Bais, H. P., Weir, T. L., Perry, L. G., Gilroy, S., & Vivanco, J. M. (2006). The role of root exudates in rhizosphere interactions with plants and other organisms. *Annual Review of Plant Biology*, 57, 233–266.

Bano, A., & Fatima, M. (2009). Salt tolerance in *Zea mays* (L.) following inoculation with *Rhizobium* and *Pseudomonas*. *Biology and Fertility of Soils*, 45(4), 405–413.

Bashan, Y., de-Bashan, L. E., Prabhu, S. R., & Hernandez, J. P. (2014). Advances in plant growth-promoting bacterial inoculant technology: formulations and practical perspectives (1998–2013). *Plant and Soil*, 378(1–2), 1–33.

Bharti, N., Pandey, S. S., Barnawal, D., Patel, V. K., & Kalra, A. (2016). Plant growth promoting rhizobacteria *Dietzia natronolimnaea* modulates the expression of stress responsive genes providing protection of wheat from salinity stress. *Scientific Reports*, 6, 34768.

Bhattacharyya, P. N., & Jha, D. K. (2012). Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture. *World Journal of Microbiology and Biotechnology*, 28(4), 1327–1350.

Bhattacharyya, R., Ghosh, B. N., Mishra, P. K., Mandal, B., Rao, C. S., Sarkar, D., Das, K., Anil, K. S., Lalitha, M., Hati, K. M., & Franzluebbers, A. J. (2015). Soil degradation in India: Challenges and potential solutions. *Sustainability*, 7(4), 3528–3570.

Bijay-Singh, & Craswell, E. (2021). Fertilizers and nitrate pollution of surface and ground water: an increasingly pervasive global problem. *SN Applied Sciences*, 3(4), 518.

Birthal, P. S., Negi, D. S., Khan, M. T., & Agarwal, S. (2015). Is Indian agriculture becoming resilient to droughts? Evidence from rice production systems. *Food Policy*, 56, 1–12.

Boddey, R. M., Urquiaga, S., Alves, B. J., & Reis, V. (2003). Endophytic nitrogen fixation in sugarcane: present knowledge and future applications. *Plant and Soil*, 252(1), 139–149.

Bulgarelli, D., Schlaeppi, K., Spaepen, S., Ver Loren van Themaat, E., & Schulze-Lefert, P. (2013). Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant Biology*, 64, 807–838.

Calderon, R. B., & Dangi, S. R. (2024). Arbuscular mycorrhizal fungi and *Rhizobium* improve nutrient uptake and microbial diversity relative to dryland site-specific soil conditions. *Microorganisms*, 12(4), 667.

Chaudhary, S., Dheri, G. S., & Brar, B. S. (2017). Long-term effects of NPK fertilizers and organic manures on carbon stabilization and management index under rice-wheat cropping system. *Soil and Tillage Research*, 166, 59–66.

Compant, S., Duffy, B., Nowak, J., Clément, C., & Barka, E. A. (2005). Use of plant growth-promoting bacteria for biocontrol of plant diseases: principles, mechanisms of action, and future prospects. *Applied and Environmental Microbiology*, 71(9), 4951–4959.

Das, A., Patel, D. P., Kumar, M., Ramkrushna, G. I., Mukherjee, A., Layek, J., Ngachan, S. V., & Buragohain, J. (2017). Impact of seven years of organic farming on soil and produce quality and crop yields in eastern Himalayas, India. *Agriculture, Ecosystems & Environment*, 236, 142–153.

Dodd, I. C., & Pérez-Alfocea, F. (2012). Microbial amelioration of crop salinity stress. *Journal of Experimental Botany*, 63(9), 3415–3428.

Druzhinina, I. S., Seidl-Seiboth, V., Herrera-Estrella, A., Horwitz, B. A., Kenerley, C. M., Monte, E., Mukherjee, P. K., Zeilinger, S., Grigoriev, I. V., & Kubicek, C. P. (2011). *Trichoderma*: the genomics of opportunistic success. *Nature Reviews Microbiology*, 9(10), 749–759.

Glick, B. R. (2010). Using soil bacteria to facilitate phytoremediation. *Biotechnology Advances*, 28(3), 367–374.

Glick, B. R. (2012). Plant growth-promoting bacteria: mechanisms and applications. *Scientifica*, 2012, 963401.

Glick, B. R. (2014). Bacteria with ACC deaminase can promote plant growth and help to feed the world. *Microbiological Research*, 169(1), 30–39.

Glick, B. R., Todorovic, B., Czarny, J., Cheng, Z., Duan, J., & McConkey, B. (2007). Promotion of plant growth by bacterial ACC deaminase. *Critical Reviews in Plant Sciences*, 26(5–6), 227–242.

Gupta, R., Anand, G., Gaur, R., & Yadav, D. (2021). Plant–microbiome interactions for sustainable agriculture: a review. *Physiology and Molecular Biology of Plants*, 27(1), 165–179.

Gupta, S., Schillaci, M., Walker, R., Smith, P. M., Watt, M., & Roessner, U. (2020). Alleviation of salinity stress in plants by endophytic plant-fungal symbiosis: Current knowledge, perspectives and future directions. *Plant and Soil*, 461(1–2), 219–244.

Gururani, M. A., Upadhyaya, C. P., Baskar, V., Venkatesh, J., Nookaraju, A., & Park, S. W. (2013). Plant growth-promoting rhizobacteria enhance abiotic stress tolerance in *Solanum tuberosum* through inducing changes in the expression of ROS-scavenging enzymes and improved photosynthetic performance. *Journal of Plant Growth Regulation*, 32(2), 245–258.

Haas, D., & Défago, G. (2005). Biological control of soil-borne pathogens by fluorescent pseudomonads. *Nature Reviews Microbiology*, 3(4), 307–319.

Harman, G. E., Howell, C. R., Viterbo, A., Chet, I., & Lorito, M. (2004). *Trichoderma* species—opportunistic, avirulent plant symbionts. *Nature Reviews Microbiology*, 2(1), 43–56.

Herridge, D. F., Peoples, M. B., & Boddey, R. M. (2008). Global inputs of biological nitrogen fixation in agricultural systems. *Plant and Soil*, 311(1–2), 1–18.

Herrmann, L., & Lesueur, D. (2013). Challenges of formulation and quality of biofertilizers for successful inoculation. *Applied Microbiology and Biotechnology*, 97(20), 8859–8873.

Kaushal, M., & Wani, S. P. (2016). Plant-growth-promoting rhizobacteria: drought stress alleviators to ameliorate crop production in drylands. *Annals of Microbiology*, 66(1), 35–42.

Kloepper, J. W., Leong, J., Teintze, M., & Schroth, M. N. (1980). Enhanced plant growth by siderophores produced by plant growth-promoting rhizobacteria. *Nature*, 286(5776), 885–886.

Köhl, J., Kolnaar, R., & Ravensberg, W. J. (2019). Mode of action of microbial biological control agents against plant diseases: relevance beyond efficacy. *Frontiers in Plant Science*, 10, 845.

Kumar, R., Gaind, S., & Nain, L. (2008). Evaluation of thermophilic fungal consortium for paddy straw composting. *Biodegradation*, 19(3), 395–402.

Kumar, S., Diksha, Sindhu, S. S., & Kumar, R. (2021). Biofertilizers: An ecofriendly technology for nutrient recycling and environmental sustainability. *Current Research in Microbial Sciences*, 3, 100094.

Kumar, S. P., Singh, N., Kishore, A., et al. (2024). Effect of integrated nutrient management on growth and yield of green gram [*Vigna radiata* (L) Wilczek]. *Journal of Food Legumes*, 37(1), 114–116.

Ladha, J. K., Dawe, D., Pathak, H., Padre, A. T., Yadav, R. L., Singh, B., Singh, Y., Singh, Y., Singh, P., Kundu, A. L., Sakal, R., Ram, N., Regmi, A. P., Gami, S. K., Bhandari, A. L., Amin, R., Yadav, C. R., Bhattarai, E. M., Das, S., ... Hobbs, P. R. (2003). How extensive are yield declines in long-term rice–wheat experiments in Asia? *Field Crops Research*, 81(2–3), 159–180.

Lehmann, A., Veresoglou, S. D., Leifheit, E. F., & Rillig, M. C. (2014). Arbuscular mycorrhizal influence on zinc nutrition in crop plants—a meta-analysis. *Soil Biology and Biochemistry*, 69, 123–131.

Mader, P., Kaiser, F., Adholeya, A., Singh, R., Uppal, H. S., Sharma, A. K., Srivastava, R., Sahai, V., Aragno, M., Wiemken, A., Johri, B. N., & Fried, P. M. (2011). Inoculation of root microorganisms for sustainable wheat–rice and wheat–black gram rotations in India. *Soil Biology and Biochemistry*, 43(3), 609–619.

Malusá, E., Sas-Paszt, L., & Ciesielska, J. (2012). Technologies for beneficial microorganisms inocula used as biofertilizers. *The Scientific World Journal*, 2012, 491206.

Meena, V. S., Maurya, B. R., & Verma, J. P. (2014). Does a rhizospheric microorganism enhance K^+ availability in agricultural soils? *Microbiological Research*, 169(5–6), 337–347.

Mortimer, P. E., Pérez-Fernández, M. A., & Valentine, A. J. (2008). The role of arbuscular mycorrhizal colonization in the carbon and nutrient economy of the tripartite symbiosis with nodulated *Phaseolus vulgaris*. *Soil Biology and Biochemistry*, 40(5), 1019–1027.

Mugabo, J. P., Bhople, B. S., Kumar, A. K., Singh, V. K., Mahmoud, E. A., Ullah, F., Elansary, H. O., & Singh, O. (2024). Insights into the tripartite interaction: effects of Arbuscular mycorrhizae and *Rhizobium* on root morphology, soil enzymes, and biochemical properties in pea cultivation in alluvial soils of Punjab, India. *Cogent Food & Agriculture*, 10(1), 2366384.

Negi, R., Yadav, N., & Yadav, A. N. (2025). Microbial biofertilizers: A paradigm shift towards agricultural sustainability. *Biologia*, 80, 1–24.

O'Callaghan, M. (2016). Microbial inoculation of seed for improved crop performance: issues and opportunities. *Applied Microbiology and Biotechnology*, 100(13), 5729–5746.

Oldroyd, G. E. (2013). Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. *Nature Reviews Microbiology*, 11(4), 252–263.

Ongena, M., & Jacques, P. (2008). *Bacillus* lipopeptides: versatile weapons for plant disease biocontrol. *Trends in Microbiology*, 16(3), 115–125.

Pandey, A., Trivedi, P., Kumar, B., & Palni, L. M. S. (2006). Characterization of a phosphate solubilizing and antagonistic strain of *Pseudomonas putida* (B0) isolated from a sub-alpine location in the Indian Central Himalaya. *Current Microbiology*, 53(2), 102–107.

Parniske, M. (2008). Arbuscular mycorrhiza: the mother of plant root endosymbioses. *Nature Reviews Microbiology*, 6(10), 763–775.

Patten, C. L., & Glick, B. R. (2002). Role of *Pseudomonas putida* indoleacetic acid in development of the host plant root system. *Applied and Environmental Microbiology*, 68(8), 3795–3801.

Pérez-Jaramillo, J. E., Mendes, R., & Raaijmakers, J. M. (2016). Impact of plant domestication on rhizosphere microbiome assembly and functions. *Plant Molecular Biology*, 90(6), 635–644.

Pieterse, C. M., Zamioudis, C., Berendsen, R. L., Weller, D. M., Van Wees, S. C., & Bakker, P. A. (2014). Induced systemic resistance by beneficial microbes. *Annual Review of Phytopathology*, 52, 347–375.

Pingali, P. L. (2012). Green Revolution: Impacts, limits, and the path ahead. *Proceedings of the National Academy of Sciences*, 109(31), 12302–12308.

Ramesh, A., Sharma, S. K., Sharma, M. P., Yadav, N., & Joshi, O. P. (2014). Inoculation of zinc solubilizing *Bacillus aryabhattai* strains for improved growth, mobilization and biofortification of zinc in soybean and wheat cultivated in Vertisols of central India. *Applied Soil Ecology*, 73, 87–96.

Richardson, A. E., & Simpson, R. J. (2011). Soil microorganisms mediating phosphorus availability update on microbial phosphorus. *Plant Physiology*, 156(3), 989–996.

Rillig, M. C. (2004). Arbuscular mycorrhizae, glomalin, and soil aggregation. *Canadian Journal of Soil Science*, 84(4), 355–363.

Rodríguez, H., & Fraga, R. (1999). Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnology Advances*, 17(4–5), 319–339.

Saha, M., Sarkar, S., Sarkar, B., Sharma, B. K., Bhattacharjee, S., & Tribedi, P. (2016). Microbial siderophores and their potential applications: a review. *Environmental Science and*

Pollution Research, 23(5), 3984–3999.

Sahil, R., Pal, V., Kharat, A. S., & Jain, M. (2025). A multi-omics meta-analysis of rhizosphere microbiome reveals growth-promoting marker bacteria at different stages of legume development. *Plant, Cell & Environment*, 48(5), 3120–3138.

Sandhya, V., Ali, S. Z., Grover, M., Reddy, G., & Venkateswarlu, B. (2009). Alleviation of drought stress effects in sunflower seedlings by the exopolysaccharides producing *Pseudomonas putida* strain GAP-P45. *Biology and Fertility of Soils*, 46(1), 17–26.

Santi, C., Bogusz, D., & Franche, C. (2013). Biological nitrogen fixation in non-legume plants. *Annals of Botany*, 111(5), 743–767.

Sapkota, T. B., & Bijay-Singh. (2025). *Roadmap for Atmanirbharta in Fertilizers* (Policy Paper 150). National Academy of Agricultural Sciences (NAAS), New Delhi.

Sasse, J., Martinoia, E., & Northen, T. (2018). Feed your friends: do plant exudates shape the root microbiome? *Trends in Plant Science*, 23(1), 25–41.

Schoebitz, M., López, M. D., & Roldán, A. (2013). Bioencapsulation of microbial inoculants for better soil–plant fertilization: a review. *Agronomy for Sustainable Development*, 33(4), 751–765.

Seneviratne, G., Zavahir, J. S., Bandara, W. M. M. S., & Weerasekara, M. L. M. A. W. (2008). Fungal–bacterial biofilms: their development for novel biotechnological applications. *World Journal of Microbiology and Biotechnology*, 24(6), 739–743.

Sessitsch, A., Pfaffenbichler, N., & Mitter, B. (2019). Microbiome applications from lab to field: facing complexity. *Trends in Plant Science*, 24(3), 194–198.

Sharma, D. K., Singh, A., Sharma, P. C., Dagar, J. C., & Chaudhari, S. K. (2016). Sustainable management of sodic soils for crop production: opportunities and challenges. *Journal of Soil Salinity and Water Quality*, 8(2), 109–130.

Sharma, T. (2024). Antibiotic and antifungal characteristics of moringa (*Moringa oleifera*). *The Journal of Rural Advancement*, 12(1), 34–49.

Singh, D., & Prasanna, R. (2020). Potential of microbes in the biofortification of Zn and Fe in dietary food grains: A review. *Agronomy for Sustainable Development*, 40(2), 15.

Singh, J. S., Koushal, S., Kumar, A., Vimal, S. R., & Gupta, V. K. (2016). Book review: Microbial inoculants in sustainable agricultural productivity—Vol. II: Functional application. *Frontiers in Microbiology*, 7, 2105.

Smith, J., Yeluripati, J., Smith, P., & Nayak, D. R. (2020). Potential yield challenges to scale-up of zero budget natural farming. *Nature Sustainability*, 3(3), 247–252.

Smith, S. E., & Read, D. J. (2008). *Mycorrhizal Symbiosis* (3rd ed.). Academic Press.

Spaepen, S., Vanderleyden, J., & Remans, R. (2007). Indole-3-acetic acid in microbial and microorganism-plant signaling. *FEMS Microbiology Reviews*, 31(4), 425–448.

Teja, M. S., Kishore, A., Sharma, J. D., Kumar, S. P., Reddy, K. S., & Pramod, T. (2022). Different levels of phosphorus and phosphorus solubilizing bacteria (PSB) influence growth, yield and economics of green gram [*Vigna radiata* L.]. *Journal of Food Legumes*, 35(3), 180–183.

Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R., & Polasky, S. (2002). Agricultural sustainability and intensive production practices. *Nature*, 418(6898), 671–677.

Tyśkiewicz, R., Nowak, A., Ozimek, E., & Jaroszuk-Ściseł, J. (2022). *Trichoderma*: The current status of its application in agriculture for the biocontrol of fungal phytopathogens and stimulation of plant growth. *International Journal of Molecular Sciences*, 23(4), 2329.

Vaidyanathan, S. (2022). *State of Biofertilizers and Organic Fertilizers in India*. Centre for Science and Environment (CSE), New Delhi.

Van Wees, S. C., Van der Ent, S., & Pieterse, C. M. (2008). Plant immune responses triggered by beneficial microbes. *Current Opinion in Plant Biology*, 11(4), 443–448.

Vandenkoornhuyse, P., Quaiser, A., Duhamel, M., Le Van, A., & Dufresne, A. (2015). The importance of the microbiome of the plant holobiont. *New Phytologist*, 206(4), 1196–1206.

Venkateswarlu, B., & Shanker, A. K. (2009). Climate change and agriculture: adaptation and mitigation strategies. *Indian Journal of Agronomy*, 54(2), 226–230.

Vessey, J. K. (2003). Plant growth promoting rhizobacteria as biofertilizers. *Plant and Soil*, 255(2), 571–586.

Vorholt, J. A., Vogel, C., Carlström, C. I., & Müller, D. B. (2017). Establishing causality: opportunities of synthetic communities for plant microbiome research. *Cell Host & Microbe*, 22(2), 142–155.

Yang, J., Kloepper, J. W., & Ryu, C. M. (2009). Rhizosphere bacteria help plants tolerate abiotic stress. *Trends in Plant Science*, 14(1), 1–4.

Zhang, H., Kim, M. S., Sun, Y., Dowd, S. E., Shi, H., & Paré, P. W. (2008). Soil bacteria confer plant salt tolerance by tissue-specific regulation of the sodium transporter *HKT1*. *Molecular Plant-Microbe Interactions*, 21(6), 737–744.

Journal of Rural Advancement
[ISSN 2347-2561 (P); 2583-6102 (E)]
Vol. 12, Issue 02, October 2024

Leveraging AI for Preservation and Promotion of Braj Folk Music: Cultural, Social, and Economic Implications for Rural Advancement

Ranu Verma

Dau Dayal Mahila Postgraduate College, Firozabad-283203 (Uttar Pradesh), India
Email: drrenuvermamusic@rediff.com
Received on: July 26, 2024; Accepted on: August 21, 2024

Abstract

The integration of Artificial Intelligence (AI) in preserving and promoting Braj folk music offers promising avenues for cultural preservation, economic growth, and rural development. This review explores how AI can enhance the documentation, classification, and accessibility of Braj folk music, ensuring its relevance in the digital age. By utilizing AI for tasks such as digitization, pattern recognition, and music composition, this research highlights the role of AI in bridging the gap between traditional cultural expressions and modern technology. However, challenges like data quality, cultural sensitivity, and ethical concerns remain significant barriers to widespread AI adoption in this domain. The paper emphasizes the need for collaborative efforts between AI experts, musicologists, and cultural custodians to ensure the preservation of the emotional and cultural depth of Braj music. Moreover, future research should examine the socio-economic impact of AI-driven music promotion on rural communities, which could benefit from increased visibility and commercialization of local art forms. This review aims to contribute to discussions on AI's role in preserving intangible cultural heritage while promoting rural economic advancement.

Keywords: Artificial Intelligence, Braj Folk Music, Machine Learning, Audio Recognition, Cultural Preservation, Musicology.

Introduction

Braj folk music is a significant cultural and spiritual heritage of the Braj region of India (Sinha, 2014). However, like many other traditional art forms, Braj folk music faces threats due to poor documentation, modernization, and declining interest among younger generations. In recent years, Artificial Intelligence (AI) has emerged as a powerful tool in musicology, providing new opportunities for the digitization and revival of traditional music (Koh, 2018).

The aim of this research paper is to review current literature on the role of AI in musicology and explore its potential applications in Braj folk music. This study examines the role of AI in digitizing, analyzing, and promoting Braj folk music through techniques like machine learning (ML), natural language processing (NLP), and automated sound recognition.

Materials and Methods

In this review, the research methodology involves an in-depth analysis of existing scientific papers and articles on the use of AI in musicology. Relevant and scientifically significant studies related to traditional music genres were selected. The key AI research areas in this context include:

1. **Audio Recognition and Classification**: Training AI models to recognize the distinct tunes, rhythms, and instruments of Braj music.
2. **Music Composition and Reproduction**: AI tools capable of creating or reproducing new music based on the patterns and rules of Braj music.
3. **Music Recommendation and Dissemination**: Using AI to reach wider audiences for Braj folk music through digital platforms.
4. **Natural Language Processing for Song Analysis**: Analyzing and studying the lyrical content of Braj songs using NLP techniques.

Results

The application of Artificial Intelligence (AI) in the field of music has ushered in transformative changes, especially in the preservation, analysis, and generation of culturally significant music. Braj folk music, an intrinsic part of Indian heritage (Slawek, 1986), stands to benefit from these advancements. Through the integration of AI, various aspects of Braj folk music, from digitization to composition, are enhanced, providing both technical and cultural solutions for its preservation and growth. The detailed scope of AI's role in this domain extends beyond initial successes, revealing opportunities for innovation, access, and global dissemination.

1. Preservation through Digitization

The preservation of Braj folk music through digitization (Gupta and Sharma, 2017) is one of the most vital contributions of AI. With machine learning (ML) models trained on distinctive patterns of Braj music, AI can assist in cataloging, archiving, and safeguarding recordings from potential degradation. Beyond simple digitization, these models offer a unique advantage by recognizing and classifying intricate aspects of Braj music, such as its specific melodic structures, rhythmic patterns, and instrumental arrangements. AI-driven technologies enable the creation of extensive digital libraries, facilitating both long-term preservation and enhanced access for future generations. The automatic metadata generation further streamlines the cataloging process, making it easier for researchers and enthusiasts to locate specific types of songs. Moreover, digital preservation ensures that Braj folk music, an integral part of India's intangible cultural heritage, remains accessible despite the inevitable wear and tear of physical recordings.

2. Pattern Recognition and Classification

The capability of AI in identifying patterns and performing classification tasks is a game changer for music research. For Braj folk music, AI algorithms can be trained to recognize recurring motifs, themes, or

instrument combinations that define the genre (Dutta et al., 2017). Such pattern recognition not only facilitates the study of musical characteristics but also aids in the classification of songs based on parameters like their thematic content (devotional, pastoral, romantic), instruments used, or song type (bhajan, kirtan, or seasonal songs). These classifications enable efficient sorting and retrieval of Braj folk songs, making AI an indispensable tool for researchers, musicologists, and even casual listeners who seek specific content within vast music libraries. The importance of accurate classification cannot be overstated, as it forms the foundation for more nuanced studies of regional musical traditions, contributing to a better understanding of cultural evolution through music.

3. Restoration of Recordings

Restoration of old and poor-quality recordings is one of the areas where AI has made tremendous strides. Many Braj folk music recordings, which are often decades old, suffer from noise, distortion, and degradation. AI-powered tools are increasingly being employed to clean, enhance, and restore these recordings. Through deep learning algorithms, these tools can isolate unwanted noise, enhance vocal clarity, and improve the overall fidelity of the music, thus reviving what might otherwise be lost to history (Hanslip, 2023). Importantly, AI restoration tools can now work with minimal human intervention, reducing the time and effort required for manual restoration processes. This enhancement not only makes the music more enjoyable but also ensures that future generations have access to these cultural artifacts in high-quality formats. The ethical considerations in this context involve maintaining the original integrity of the music while improving its accessibility.

4. Cultural and Lyrical Analysis

AI's ability to conduct Natural Language Processing (NLP) has significant implications for the study of Braj folk music (Obaid, et al., 2023). The lyrics of these songs are deeply rooted in historical, cultural, and religious contexts, and AI-driven NLP models can analyze lyrical content to extract meaningful insights. Such analyses can reveal underlying themes, cultural values, and historical events referenced in the songs, providing a richer understanding of the societal context in which the music was created. For instance, sentiment analysis could uncover emotional trends in certain types of songs, while topic modeling could help categorize songs by thematic relevance to historical events, folklore, or religious practices. Moreover, AI can uncover subtle linguistic patterns that might be lost in human analysis, such as recurring metaphors, shifts in language over time, or regional variations in dialect. These insights can further inform anthropological studies, helping researchers understand how Braj folk music reflects broader cultural shifts.

5. Music Composition

The creative potential of AI in music composition has expanded rapidly in recent years (Anantrasirichai and Bull, 2022), and although in its nascent stages, it holds promise for Braj folk music. AI models trained on traditional Braj musical structures can generate new compositions that adhere to the rules and styles characteristic of this genre. These generative AI models can emulate traditional melodic scales, rhythmic patterns, and harmonic structures, potentially reviving interest in the genre by creating new, yet familiar, compositions. This AI-generated music could serve various purposes, from education and cultural promotion to the reinvigoration of contemporary music styles with traditional

roots. Additionally, AI's generative capabilities could allow composers to experiment with fusions of Braj folk music with other genres, fostering innovation while preserving core musical traditions.

6. Enhancing Popularity on Digital Platforms

The digital age has brought unprecedented opportunities for music to reach global audiences, and AI plays a central role in making this possible. Recommendation systems driven by AI are widely used on streaming platforms and can help Braj folk music gain visibility among diverse audiences. By analyzing user preferences and listening habits, AI can recommend Braj folk songs to listeners who may not otherwise encounter this genre, potentially broadening its appeal. Furthermore, AI-powered analytics can help musicians and cultural organizations understand the demographic and geographic distribution of listeners, providing valuable data for targeted promotion strategies. These recommendation systems ensure that Braj folk music maintains its relevance in an increasingly digital world (Gupta, 2011), connecting traditional music with contemporary global audiences.

7. Reviving Folk Traditions through Collaboration

AI offers a unique opportunity to not only preserve but also revitalize traditional folk music through collaboration with human musicians (Yang and Huang, 2023). Interactive AI systems allow musicians to collaborate with AI models to create innovative compositions or reinterpret traditional works. In the context of Braj folk music, this could mean reimagining classical compositions with modern instrumentation or creating fusion pieces that bring Braj music into new genres, such as electronic or world music. Such collaborations can serve as a bridge between past and present, maintaining the cultural relevance of Braj music while simultaneously adapting to contemporary tastes. Moreover, the use of AI in music education can promote the learning of Braj folk music among younger generations, ensuring that these traditional forms continue to thrive in the future.

8. Cross-cultural Exchange and Research

Another significant impact of AI is its facilitation of cross-cultural exchange in music (Shadiev, 2019). AI technologies enable the analysis of Braj folk music within the broader context of global folk traditions, allowing for comparative studies between different cultures. This not only enhances academic research but also opens doors for intercultural musical exchanges, where artists from different traditions can draw inspiration from Braj folk music and incorporate its elements into their own work. AI's ability to identify commonalities and differences between musical forms contributes to the global appreciation of regional music like Braj folk, elevating it as a subject of international interest and scholarly inquiry.

Discussion

The integration of Artificial Intelligence (AI) in preserving and promoting Braj folk music is not merely confined to traditional preservation techniques. By harnessing AI, researchers, ethnomusicologists, and musicians can explore traditional music through advanced analytical methods, uncovering deeper insights into the intricacies of the genre. AI allows the analysis of regional variations within Braj folk music, which can reveal the subtle differences in style, tempo, and instrumentation that define distinct local traditions. Such insights would be difficult to attain using conventional musicological

techniques but are made possible by AI's ability to process and identify complex patterns within vast datasets.

For instance, AI-based musicological tools can map the evolution of Braj folk music over decades (Coppolillo et al., 2023), illustrating how various socio-cultural influences have shaped its development. This level of analysis may identify significant trends in musical styles, such as shifts in lyrical themes, the introduction of new instruments, or changes in vocal techniques. This kind of AI-driven musicological research could offer novel insights into the ways that Braj folk music has adapted over time to external influences like migration, technological advancements, or changes in the region's social fabric. The analysis of such historical trends can be particularly valuable in understanding how traditional art forms evolve while still maintaining their cultural essence.

In addition, AI introduces new opportunities for the promotion of Braj folk music on digital platforms. Streaming services equipped with AI-driven recommendation systems can help ensure the visibility of this genre by suggesting it to listeners who may not be familiar with it but are likely to appreciate its stylistic elements. These systems, powered by algorithms that learn user preferences, can help bring niche genres like Braj folk music to a broader, global audience. AI can analyze user listening patterns to suggest Braj music to people with similar tastes, thus expanding its reach far beyond local communities. This modern form of digital marketing holds tremendous potential for ensuring the relevance of Braj folk music in an increasingly globalized and digital music industry.

However, despite the immense potential, the implementation of AI in this field comes with its set of limitations. One of the critical issues lies in the quality of datasets required for machine learning algorithms to function effectively. AI models rely heavily on comprehensive, high-quality training data to perform accurately. In the case of Braj folk music, many existing recordings are of poor quality, fragmented, or incomplete, which makes it challenging to train models that can accurately process and replicate the intricacies of the music. Older recordings, often the most authentic representations of Braj music, may have been recorded on outdated equipment, resulting in substandard audio quality. While AI can clean and restore some of these recordings, the success of these efforts largely depends on the initial quality of the audio data, presenting a significant hurdle for researchers and archivists.

Challenges and Limitations

AI applications in Braj folk music also raise several important challenges, particularly related to cultural sensitivity, data quality, and ethical considerations.

1. Lack of High-Quality Data

The availability and quality of training data significantly impact the success of AI models. Machine learning and deep learning models require vast amounts of well-annotated, high-quality audio recordings to effectively "learn" and reproduce musical elements. In the context of Braj folk music, much of the available material might be poorly preserved, incomplete, or recorded with inadequate technology. This presents a barrier to the effective application of AI, as models trained on low-quality data are less likely to accurately capture the true essence of the music. Research in AI emphasizes that the "garbage in, garbage out" principle applies particularly strongly in this area, where low-quality input data will yield suboptimal results from AI models. Moreover, the fragmented nature of the

available datasets might lead to incomplete representations of the musical genre, limiting the scope of AI-driven analyses.

Efforts to create a more robust dataset would involve not only digitizing and restoring old recordings but also commissioning new, high-quality recordings of Braj folk music. This could serve as a critical resource for AI training, ensuring more accurate and comprehensive models. In doing so, collaboration between AI experts, ethnomusicologists, and traditional musicians is essential to ensure that the data being used is both authentic and of high quality.

2. Cultural Sensitivity

AI has a known limitation when it comes to interpreting and reproducing cultural contexts (Elish, 2017). While machine learning models excel at recognizing patterns and replicating them, they are often unable to grasp the deeper cultural, spiritual, and emotional nuances that are intrinsic to traditional music forms like Braj folk music. AI-generated compositions might accurately replicate the melody or rhythm of Braj music, but they may lack the emotional depth and authenticity that comes from human interpretation and cultural experience. The spirituality, storytelling, and communal aspects of Braj music, which often reflect centuries of religious and cultural practices, cannot easily be reproduced by an algorithm. Research has shown that while AI can simulate the technical components of music, it struggles with the more abstract, emotional qualities that are often central to traditional music genres.

Additionally, the process of distilling Braj folk music into patterns for AI to analyze risks reducing the cultural significance of the music to a series of mathematical relationships. This is a challenge that extends beyond just Braj music; ethnomusicologists have long cautioned against the over-reduction of traditional and indigenous music forms into technical components without considering their cultural context. It is crucial that AI applications in Braj folk music be developed in consultation with cultural experts and musicians to ensure that the spiritual and emotional elements of the music are not lost in the process.

3. Ethical Considerations

The use of AI in music creation also brings forth important ethical considerations, particularly around issues of authorship, ownership, and cultural appropriation. As AI-generated music becomes more common, questions arise about who holds the rights to these compositions (Bonadio and McDonagh, 2018). In the case of Braj folk music, which is deeply rooted in community practices, it is essential to ensure that the original creators and performers receive appropriate recognition and compensation for their work. There is a risk that AI-generated music could be commodified without giving due credit to the artists and communities from which the music originated. Research in the field of AI ethics stresses the importance of ensuring that intellectual property rights are upheld and that traditional artists are not sidelined in the pursuit of technological advancement.

Moreover, the commercialization of AI-generated Braj music could lead to a form of cultural exploitation, where the music is used for profit without regard to its cultural and spiritual significance. AI-generated compositions could be appropriated by commercial interests in ways that strip the music of its original meaning, further exacerbating issues of cultural misrepresentation and exploitation. Thus, it is critical to establish ethical guidelines and frameworks that protect the cultural

ownership of Braj folk music and ensure that any AI-generated music is respectful of its origins.

Future Directions

Future research in AI-driven preservation and promotion of Braj folk music must address several key areas (Pauwels, 2021). First, there is a pressing need to create comprehensive, high-quality datasets of Braj folk music. The scarcity of well-documented and digitized recordings currently limits the effectiveness of AI models. Developing an expansive archive that includes various styles, instruments, regional variations, and historical recordings would greatly enhance AI's ability to accurately analyze and reproduce Braj folk music. These datasets should also include metadata on cultural context, historical significance, and lyrical meaning to improve the richness and usability of AI-generated insights.

Another area of focus should be on enhancing the cultural competency of AI models. AI systems must not merely recognize musical patterns but must also capture the emotional depth, cultural nuances, and spiritual essence that are integral to Braj folk music. This can be achieved through interdisciplinary collaboration between AI researchers, musicologists, and cultural experts. Training AI models with datasets that reflect the subtleties of this genre would allow for more authentic reproduction and analysis, while preventing the homogenization of this rich cultural heritage.

Moreover, future studies should explore how AI-driven music promotion impacts both the social and economic ecosystems of rural communities in Braj. While AI can amplify the visibility of Braj folk music on global platforms, it is important to assess how this newfound exposure benefits the local artists and musicians. Research should examine whether AI-enabled global exposure translates into sustainable income and enhanced social standing for these communities. Such studies should also explore the risks of commercialization, such as the potential loss of cultural authenticity or exploitation of traditional artists.

Furthermore, the long-term impact of AI interventions on cultural ecosystems needs to be analyzed. AI's ability to influence musical tastes and reshape how regional music is consumed globally could lead to changes in the way local communities engage with their own cultural practices. Studies that investigate the role of AI in shaping the future of cultural traditions will be essential to ensure that technological advancement does not lead to cultural erosion.

Finally, future research must explore the ethical dimensions of AI-driven music creation and promotion. Issues such as intellectual property rights, cultural appropriation, and fair compensation for local artists should be at the forefront of discussions. By addressing these challenges, AI can not only serve as a tool for cultural preservation but also ensure that the communities whose music it promotes are equitably represented and rewarded.

Conclusion

AI's potential in preserving Braj folk music offers a critical tool for both cultural and rural development. This paper has highlighted the immense possibilities AI presents for the preservation, analysis, and promotion of Braj folk music, ensuring its survival and continued relevance in the digital era. However, challenges such as data quality, cultural sensitivity, and ethical concerns must be addressed through multidisciplinary collaboration. AI initiatives should incorporate the expertise of cultural custodians, ethnomusicologists, and rural

communities to ensure that the emotional and cultural depth of Braj music is preserved. Furthermore, the socio-economic impact of promoting regional music through AI needs to be thoroughly examined to ensure that rural communities benefit from the global recognition and commercialization of their cultural heritage. Ultimately, AI can serve as a powerful tool in both preserving intangible cultural heritage and contributing to rural advancement, provided that the cultural and ethical dimensions are carefully managed.

References

Anantrasirichai N and Bull D. 2022. Artificial intelligence in the creative industries: A review. Artificial Intelligence Review **55**:589–656.

Bonadio E and McDonagh L. 2018. Artificial intelligence as producer and consumer of copyright works: Evaluating the consequences of algorithmic creativity.

Coppolillo E, Guarascio M, Minici M and Pisani F S. 2023. Exploiting deep learning and explanation methods for movie tag prediction. In IDEAS '23: Proceedings of the 27th International Database Engineered Applications Symposium, pp 177-184.

Datta A K, Solanki S S, Sengupta R, Chakraborty S, Mahto K and Patranabis A. 2017. Signal analysis of Hindustani classical music. Springer.

Elish M C and Boyd D. 2017. Situating methods in the magic of Big Data and AI. Communication Monographs **85**(1):57–80.

Gupta B M. 2011. Science and folk music. Indian Journal of Science Communication **10**(2):23-30.

Gupta D K and Sharma V. 2017. Enriching and enhancing digital cultural heritage through crowd contribution. Journal of Cultural Heritage Management and Sustainable Development **7**(1):14-32.

Hanslip M. 2023. Applications of discriminative, generative and predictive deep learning processes to solo saxophone practice. Thesis, Doctor of Philosophy, University of York.

Koh A. 2018. Music for AI reports: Dual prospects in music production. Thesis, Bachelor of Arts, City University, New York.

Obaid A J, Bhushan B, Muthmainnah S and Suman R S. 2023. Advanced applications of generative AI and natural language processing models. IGI Global.

Pauwels H. 2021. Cultivating emotion and the rise of the vernacular: The role of affect in ‘early Hindi-Urdu’ song. South Asian History and Culture **12**(2–3):146–165.

Shadiev R, Sun A and Huang Y M. 2019. A study of the facilitation of cross-cultural understanding and intercultural sensitivity using speech-enabled language translation technology. British Journal of Educational Technology **50**(3):1415-1433.

Sinha A. 2014. The sacred landscape of Braj, India: Imagined, enacted, and

reclaimed. Landscape Journal **331**:59-75.

Slawek S M. 1986. Kirtan: A Study of The Sonic Manifestations of The Divine in The Popular Hindu Culture of Banaras (Ethnomusicology, Folk Music, India, Religion, South Asia). University of Illinois at Urbana-Champaign.

Yang Y and Huang X. 2023. Research based on the application and exploration of artificial intelligence in the field of traditional music. Journal of Sensors **2023**(1).

Journal of Rural Advancement
[ISSN 2347-2561 (P); 2583-6102 (E)]
Vol. 12, Issue 02, October 2024

Review of Social Media's Influence on School Management: Urban vs Rural Perspectives

Shivani Bhadauria

[1]Department of Management, Maharishi School of Commerce & Management, Maharishi University of Information Technology, Lucknow-226013 (Uttar Pradesh), India
Email: shivani7994@yahoo.in
Received on: August 1, 2024; Accepted on: August 28, 2024

Abstract

This paper provides a comprehensive review of the impact of social media on school management, particularly focusing on the distinct challenges and opportunities that urban and rural private junior high schools face. It highlights how urban schools, equipped with better technological infrastructure and resources, often adopt sophisticated, multi-platform social media strategies for communication, marketing, and crisis management. These schools utilize advanced analytics, engage diverse audiences, and implement complex content strategies across platforms like Facebook, Instagram, and Twitter. Conversely, rural schools, despite facing limitations such as slower internet speeds and fewer resources, effectively leverage their closer-knit communities. These schools often focus on a single platform, like Facebook, to build strong, personal connections with stakeholders, ensuring consistent and targeted messaging. The analysis underscores the necessity of context-specific strategies in social media use, with urban schools benefiting from scale and complexity, while rural schools excel through community-centered approaches. The paper also offers policy recommendations to enhance social media capabilities in rural schools and proposes guidelines for urban schools to manage the complexities of social media use. Additionally, it calls for future research to explore the long-term impacts of social media on educational outcomes and the potential role of emerging technologies in refining these strategies.

Keywords: Community Engagement, Crisis Management, Digital Literacy, Rural Schools, Social Media Strategies, Urban Schools.

Introduction

Social media has knowingly impacted many facets of school administration and has occurred as Social media has knowingly impacted many facets of school administration and has occurred as an essential instrument in today's educational environment. Social media offers probable and problems for private junior high schools, especially those in urban and rural settings, with regard to marketing, stakeholder engagement, and communication (Gulati, 2008). With the help of the platforms, schools may interact more dynamically with students and parents, reach a wider audience, and foster a feeling of community that knows no geographical boundaries. However, there are distinguished differences in the effectiveness of social media in school management across urban and rural environments, depending on

factors including resource availability, community structure, and technological access (Bandyopadhyay et al., 2021). This review study has been conducted to compare and contrast the ways in which social media has affected the management approaches employed by private junior high schools in these two different settings i.e. urban and rural. By being more conscious of the distinctions and similarities, school administrators can adapt their methods and practices to better suit the needs of their local communities.

Private junior high schools located in the cities frequently operate in ruthless markets where distinction through branding and communication is essential. Babu et al., (2019) reported that these educational institutions can use social media like WhatsApp and facebook etc. as a potent tool to highlight their successes, draw in new students, and interact with stakeholders instantly. The higher level of digital literacy in urban localities compared to rural and the general accessibility of high-speed internet, additionally enhance the impact of social media in urban environments. Social media platforms are being utilized by urban schools to engage their audiences in a variety of cutting-edge ways. Transparency and participation both have been replaced by efficient use of live streaming of various events, frequent updates on academic activity, and interactive meetings between school administration and parents.

On the other hand, low levels of digitally literate local population and limited access to technology frequently hinder the use of social media in school management in rural locations (Gulati, 2008). Due to weak internet signals, educational institutions in rural areas are not in position to fully explore social media sites like YouTube. The rural educational organizations can still use these platforms for sharing success stories of their students, building stronger community ties, and interacting more personally with parents and students. For illustration, social media—especially WhatsApp—has mostly been used by rural schools to highlight student accomplishments, interact with local leaders, and exclusively promote local culture. Additionally, this social media is employed to assure that parents participate more in the decision-making process of the school.

Theoretical Framework

The role of social media in management of the schools in rural communities is applied in several theoretical perspectives, like communication theories, network theories, and stakeholder engagement models (McPhail, 2010). These theories suggested that social media is capable to enhance communication competence, simplify broader outreach, and involve stakeholders in a more dynamic and interactive way. Communication theory, for an example, speculates that the involvement of social media may break down traditional barriers to information flow, enabling schools to disseminate information and communication quickly and widely (Kumar, 2020). This is mainly relevant in the context of urban compared to rural schools, where timely communication is critical for managing large and diverse students. The ability to rapidly inform a wide audience about either changes in school policy, upcoming events, or emergency situations demonstrates how social media supports efficient communication processes.

Theoretical Framework

Network theory further elucidates the capabilities of social media to create an interconnected network of stakeholders, including students, parents, teachers, administration and community members

(Henderson, 2015). In urban environments, these networks are very complex and expansive, allowing for the rapid spreading of information and fostering collaborative relationships. The schools in urban areas often have better access to advanced technological infrastructure which has the capacity to enable them for developing more complicated social media strategies, like content creation teams for producing high-quality multimedia data. These resources, ranging from promotional videos of student success stories, are disseminated quickly and broadly through interconnected networks.

Even if they operate on smaller, more localized networks, rural schools may use social media to strengthen their relationships and stand-in a feeling of community in order to their academic objectives (Moitra et al., 2015). In order to stay in touch with families of students who may live far from the school, rural schools are known to employ common social media platforms like WhatsApp. An access to social media allows the family of the students to stay updated regarding the school events and become a part of their ward's educational journey. Social media gives rural schools the opportunity to highlight their involvement in the local community and regional customs, which can make parents and students proud of them.

Stakeholder engagement models, such as Freeman's (1984) stakeholder theory, also highlight the importance of social media in managing relationships with key stakeholders. For urban schools, social media provides a platform for engaging a diverse array of stakeholders, including parents, alumni, and local businesses, in meaningful dialogue. In rural areas, where schools often play a central role in the community, social media can be used to engage stakeholders in a more personalized and direct manner, fostering a sense of shared responsibility for educational outcomes (Naidu, 2019).

Urban vs. Rural School Management

The literature highlights significant differences in the management challenges faced by urban and rural schools. Urban schools often have better access to technology, which enables more sophisticated use of social media. They can implement advanced digital marketing strategies, engage in real-time communication with a large and diverse audience, and utilize data analytics to tailor their messaging and outreach efforts. For example, urban schools can use social media platforms to conduct virtual tours, host live Q&A sessions, and share multimedia content that showcases their facilities, faculty, and student achievements.

In contrast, rural schools may struggle with technological limitations but often enjoy closer-knit communities that can be leveraged through more personalized social media strategies (Payyappilly Jose, 2020). The sense of community in rural areas often translates into stronger relationships between schools and families, which can be enhanced through social media by sharing local success stories, highlighting community events, and facilitating direct communication between parents and teachers (Gautam, 2014). Despite the challenges, rural schools can use social media to build and maintain these relationships, ensuring that all stakeholders remain informed and engaged in the educational process.

Moreover, the disparity in resources between urban and rural schools often leads to differences in how social media is integrated into school management. Urban schools, with greater access to financial and technological resources, are more likely to invest in professional social media management,

including dedicated staff or agencies that handle their online presence (Venkatraman, 2017). Rural schools, on the other hand, may rely on teachers or administrators to manage their social media accounts, which can lead to a more authentic but less polished online presence. This difference in resource allocation impacts the effectiveness of social media strategies and the extent to which schools can engage their stakeholders.

Another critical difference lies in the target audience of social media efforts. Urban schools often cater to a more diverse and transient population, which requires a broader and more flexible approach to communication (Rao, 2005). In contrast, rural schools typically serve a more stable and homogeneous population, allowing for more targeted and community-focused social media strategies. This distinction influences not only the content shared on social media but also the platforms chosen for engagement. For instance, urban schools may prioritize platforms like Instagram and Twitter, which appeal to younger, more tech-savvy audiences, while rural schools might focus on Facebook or local community networks that resonate more with their audience (Venkatraman, 2017).

Conclusion

The conclusion of this review underscores the critical role of social media in shaping the management strategies of private junior high schools, with significant differences observed between urban and rural settings. Urban schools, equipped with better technological infrastructure and resources, leverage social media to implement sophisticated, multi-platform strategies that cater to a diverse audience. These schools use social media for marketing, parent-teacher communication, crisis management, and more, often employing data-driven approaches to enhance engagement and effectiveness.

Conversely, rural schools, despite facing challenges such as limited technology access and lower social media penetration, effectively use these platforms to foster strong community ties and maintain direct, personal communication. Their simpler, more focused strategies often resonate deeply with their close-knit communities.

The review highlights the importance of context in developing social media strategies, emphasizing that urban and rural schools must tailor their approaches to their unique environments. The findings suggest that with the right support, including training and resource allocation, rural schools can enhance their social media presence. Additionally, the review calls for policies that promote equitable access to technology and encourage collaboration between urban and rural schools, ensuring that all schools can effectively utilize social media to achieve their educational and management goals.

References

Gulati S. 2008. Technology-enhanced learning in developing nations: A review. The International Review of Research in Open and Distributed Learning, **9**(1).

Bandyopadhyay S, Bardhan A, Dey P and Bhattacharyya, S. 2021. Bridging the education divide using social technologies. Springer.

Babu R, Kalaivani S and Saileela K. 2019. Empowering India Through Digital Literacy, Vol 1. Lulu. com.

McPhail T L. 2010. Global communication: Theories, stakeholders, and trends. John Wiley & Sons.

Kumar K J. 2020. Mass communication in India. Jaico publishing house.

Henderson M L. 2015. What a Tangled Wed: Exploring the Role of Social Networks in the Implementation of a Global Education Initiative. North Carolina State University.

Moitra A, Das V, Vaani G, Kumar A and Seth A. 2016. Design lessons from creating a mobile-based community media platform in Rural India. In Proceedings of the eighth international conference on information and communication technologies and development, pp 1-11.

Naidu L. 2019. An exploration of community engagement in rural India: a case study. Thesis, Doctor of Philosophy, Bhagat Phool Singh Mahila Vishwavidyalaya, Sonipat.

Payyappilly Jose S. 2020. The Nature and Structuration of Management Control in Knowledge Intensive Work A Study of Information Technology Services Work in India. Thesis, Doctor of Philosophy, University of Leicester.

Gautam M. 2014. Role and participation of parent teacher association in school management: Evidences from school of Madhya Pradesh. Madhya Pradesh Journal of Social Sciences. **19**(1):76-87.

Venkatraman S. 2017. Social media in south India. UCL press.

Rao S S. 2005. Bridging digital divide: Efforts in India. Telematics and informatics. **22**(4):361-375.

Journal of Rural Advancement
[ISSN 2347-2561 (P); 2583-6102 (E)]
Vol. 12, Issue 02, October 2024

Value Addition in Buffalo Milk for Advancement of Rural Economy

Pravin Kumar
Pandit Deen Dayal Upadhyaya Pashu Chikitsa Vigyan Vishwavidyalaya Evam Go-Anusandhan Sansthan, Mathura-281001 (Uttar Pradesh), India
Email: pravinmthr@gmail.com
Received on: September 3, 2024; Accepted on: September 25, 2024

Abstract

The present study evaluates the impact of value addition in buffalo milk production on the rural economy of Mathura District, Uttar Pradesh. By converting raw buffalo milk into value-added products such as paneer, curd, butter, ghee, and khoa, significant income enhancements were observed among rural farmers engaged in milk production. The study was conducted across five tehsils, where ten villages from two randomly selected blocks per tehsil were chosen, covering 500 farmers involved in buffalo milk production and value addition. Data collection included milk yield from different buffalo breeds, production costs, and income generation from both raw milk and processed products. The results showed a 50-70% increase in income from value-added products compared to raw milk sales. Ghee and paneer emerged as the most profitable products due to their high consumer demand and longer shelf life. The study emphasizes the potential of value addition to not only increase farmer income but also stimulate rural employment and economic growth. Policy recommendations include improving access to processing technologies, market linkages, and technical training for small-scale dairy farmers to capitalize on the benefits of value addition.

Keywords: Buffalo milk, Dairy products, Income generation, Rural economy, Value addition, Value-added products

Introduction

Buffalo milk plays an important role in India's dairy sector, contributing approximately 50% of the total milk produced nationwide (Subbanna et al., 2021). Its high fat and protein content make it suitable for producing various value-added dairy products such as paneer, khoa, curd, butter, and ghee (Khetra et al., 2022). Despite the higher demand and profitability of these products, rural buffalo farmers still primarily sell raw milk, missing out on the economic benefits of value addition (Kashyap and Bhuyan, 2023). This research paper aims to explore how the value addition of buffalo milk products can significantly enhance the rural economy by increasing farmer incomes and expanding employment opportunities.

The significance of buffalo milk in rural areas goes beyond nutrition, serving as an economic backbone for many smallholder farmers (Srivastava, 2015). The breeds studied include Murrah, Bhadawari, Mehsana, and non-descriptive buffaloes, which are prevalent in different parts of Uttar Pradesh (Muthukumar et al., 2018). Through value addition, farmers can diversify their income sources and capture a larger share of the market. The objective of this work was to compare the costs and incomes associated with raw milk and value-added products, examine consumer preferences, and evaluate the impact of value addition on the livelihoods of rural farmers.

Materials and Methods

This study was conducted in the Mathura District of Uttar Pradesh, India, which comprises five tehsils: Mathura, Chhata, Mant, Goverdhan, and Baldeo. The district is well known for its agricultural and dairy farming practices, particularly buffalo milk production. To gain a comprehensive understanding of buffalo milk value addition and its impact on rural income, a multi-stage random sampling method was employed.

Study Area Selection

In the first stage, **two blocks** were randomly selected from each of the five tehsils. This ensured adequate geographic representation across the district, capturing both densely populated areas and rural regions with varying levels of agricultural activity.

Village and Farmer Selection

From each of the randomly selected blocks, ten villages were randomly chosen, making a total of 100 villages across the study area. This random selection helped in minimizing bias and ensured that the results were representative of the broader population engaged in buffalo milk production in the district.

Within each selected village, five buffalo farmers who were actively involved in milk production, value addition (such as processing into products like ghee, paneer, curd, butter, and khoa), and marketing of these products were selected, bringing the total sample size to 500 farmers. The selection of these farmers was based on their participation in both traditional raw milk sales and value-added dairy processing. This enabled a comparative analysis of income

generation from raw milk sales and value-added products.

Data Collection

Primary data were collected using structured questionnaires and interviews with the farmers. The questionnaire covered various aspects, including:

- **Milk yield** from different buffalo breeds such as Murrah, Bhadawari, Mehsana, and non-descriptive buffaloes.
- **Production costs** for raw milk and value-added products like ghee, paneer, butter, curd, and khoa.
- **Income generation** from both raw milk and processed dairy products.
- **Consumer preferences** for buffalo milk products in local and nearby markets.
- **Challenges** faced in value addition, such as access to processing technology, storage facilities, and market access.

Secondary data were obtained from local dairy cooperatives, government reports, and academic research papers to supplement the primary data and provide context for the economic analysis.

Data Analysis

The collected data were analyzed using both descriptive and inferential statistical methods. Comparative income analysis between raw milk sales and value-added products was performed to quantify the potential benefits of value addition. Additionally, cost-benefit analysis was conducted to evaluate the profitability of producing and marketing various dairy products. The results were tabulated and interpreted in the context of rural economic advancement through buffalo milk value addition.

Results and Discussion

Comparative Income Analysis (Raw Milk vs. Value-Added Products)

Table 1 presents a comparative income analysis between selling raw buffalo milk and processing it into value-added dairy products such as paneer, curd, butter, ghee, and khoa. The table highlights the significant income difference generated by value addition. While raw milk sales yield an average income of ₹35 per liter, converting the milk into products like paneer can increase revenue to ₹150 per kilogram, while ghee commands up to ₹300 per kilogram. This suggests that value addition can increase farmers' income by over 50%, offering a more lucrative alternative to simply selling raw milk.

Value-added products have higher market demand due to their longer shelf life, higher fat content, and wider applications in Indian cuisine. For example, paneer and ghee are extensively used in daily cooking, festivals, and special occasions, which boosts their market value. Moreover, the table reflects that although processing these products requires initial investments in infrastructure and skills, the returns far outweigh the costs. It is clear that shifting focus from raw milk to value-added products can significantly enhance rural income, especially in regions

where milk production is a primary source of livelihood. The results emphasize the need for processing facilities, better market access, and technical training to enable farmers to capitalize on this profitable opportunity.

Table-1: Comparative Income Analysis (Raw Milk vs. Value-Added Products)

Product	Average Income per Liter (₹)	Value-Added Product Income (₹)
Raw Milk	35	-
Paneer	-	150
Curd	-	70
Butter	-	130
Ghee	-	300
Khoa	-	120

This analysis highlights the necessity of establishing processing units and training farmers in value addition techniques to optimize their economic returns. The value-added product market is growing in India, driven by consumer demand for quality and taste. By entering this market, rural farmers can ensure higher profits and more sustainable livelihoods.

Cost of Production for Various Buffalo Milk Products

Table 2 outlines the cost of production for various value-added buffalo milk products, including paneer, curd, butter, ghee, and khoa. This cost analysis is crucial for understanding the economics of producing these products at the farm or small-scale processing level. Paneer and ghee are shown to have relatively high production costs, with paneer costing ₹100 per kilogram and ghee costing ₹250 per kilogram. However, the market prices for these products are also high, at ₹150 and ₹300 per kilogram, respectively, making them very profitable despite their higher processing costs.

The production of curd, butter, and khoa involves lower costs, with curd costing ₹40 per kilogram, butter ₹80, and khoa ₹90. These products also have favorable market prices, making them accessible for small-scale producers. The table illustrates that even though value-added products require an investment in processing, packaging, and labor, the potential profit margins make these endeavors highly beneficial for rural farmers. By shifting their focus from raw milk sales to these higher-value products, farmers can achieve better financial stability and resilience. Additionally, producing these items locally can reduce transportation costs, improve shelf life, and increase the availability of fresh dairy products in the rural market, further enhancing profitability.

Table-2 Cost of Production for Various Buffalo Milk Products

Product	Cost of Production (₹/kg)	Market Price (₹/kg)
Paneer	100	150
Curd	40	70
Butter	80	130
Ghee	250	300
Khoa	90	120

Processing buffalo milk into these products incurs varying costs depending on the complexity of the process. However, the higher sale prices more than compensate for these costs, making value addition a viable strategy for improving rural incomes.

Milk Yield and Income from Different Buffalo Breeds

Table 3 compares the milk yield and corresponding income from various buffalo breeds, including Murrah, Bhadawari, Mehsana, and non-descriptive buffaloes. Murrah buffaloes show the highest milk yield, producing an average of 12 liters per day, which translates into an income of ₹420 per day from raw milk and ₹1500 per day from value-added products. Mehsana and Bhadawari breeds, producing 10 and 8 liters per day respectively, offer lower but still substantial incomes when their milk is processed into products such as paneer, ghee, and khoa. The non-descriptive buffaloes, which are more common in rural areas, produce only 6 liters per day and generate the lowest income both from raw milk and value-added products.

Table-3 Milk Yield and Income from Different Buffalo Breeds

Breed	Average Milk Yield (Liters/day)	Income from Raw Milk (₹)	Income from Value-Added Products (₹)
Murrah	12	420	1500
Mehsana	10	350	1200
Bhadawari	8	280	1000
Nondescriptive	6	210	850

This table emphasizes the economic advantage of investing in high-yielding breeds like Murrah and Mehsana, especially when farmers aim to maximize their income through value-added dairy production. The differences in income between raw milk sales and value-added products are stark, with value addition significantly enhancing profitability. Furthermore, the table suggests that farmers raising low-yielding, non-

descriptive buffaloes could benefit from adopting value addition to compensate for the lower milk production. This comparison underscores the importance of breed selection and milk processing in optimizing dairy farming income in rural areas.

Murrah buffaloes, due to their superior yield and fat content, offer the highest returns both in terms of raw milk and value-added products. Farmers raising nondescriptive buffaloes may benefit from focusing on higher-value products to compensate for the lower milk yield.

Consumer Preferences for Buffalo Milk Products

Table 4 provides insights into consumer preferences for different buffalo milk products, including ghee, paneer, curd, and khoa. The data reveal that ghee is the most preferred product, with 45% of consumers indicating it as their top choice. Paneer follows at 30%, curd at 15%, and khoa at 10%. This consumer preference is largely driven by the cultural and culinary significance of ghee and paneer in Indian households. Ghee is a staple in many Indian diets and is often used for cooking, religious ceremonies, and medicinal purposes. Paneer, similarly, is widely used in Indian cuisine, particularly in vegetarian dishes.

The relatively lower preference for curd and khoa suggests that while these products are also popular, they may be more regionally consumed or used in specific dishes. The table highlights the importance of focusing on products that align with consumer demand when adding value to buffalo milk. Producers aiming to maximize profits should prioritize the production of ghee and paneer, as these products have a more robust market demand. Understanding consumer preferences is key to aligning production efforts with market needs, thereby enhancing sales and profitability in rural dairy markets.

Table-4 Consumer Preferences for Buffalo Milk Products

Product	Percentage of Consumers Preferring (%)
Ghee	45
Paneer	30
Curd	15
Khoa	10

The preference for ghee and paneer reflects their widespread use in Indian cuisine and festivals. This indicates the need for farmers and small-scale dairy processors to focus on these high-demand products.

Shelf Life of Buffalo Milk Products

Table 5 examines the shelf life of various buffalo milk products, an important factor in determining the logistics of production, storage, and market distribution. Ghee has the longest shelf life at 180 days, making it the most durable product for rural producers, particularly those who lack advanced refrigeration and transportation

infrastructure. Butter has a shelf life of 30 days, which still provides flexibility in storage and sales. Paneer and khoa, on the other hand, have shorter shelf lives of 7 and 15 days respectively, which necessitates faster turnover and more efficient market distribution to prevent spoilage.

Curd, with a shelf life of 10 days, also requires prompt marketing but is easier to store than paneer or khoa due to its fermented nature. The table suggests that products like ghee and butter, with longer shelf lives, are more suitable for rural producers who wish to expand their market reach, as these products can be transported over longer distances without the risk of spoilage. Shorter shelf-life products, while still profitable, require better infrastructure, such as cold storage facilities, to maintain product quality. This analysis underscores the need for strategic planning in the production and distribution of buffalo milk products, particularly in regions where infrastructure may be underdeveloped.

Table-5 Shelf Life of Buffalo Milk Products

Product	Shelf Life (Days)
Ghee	180
Paneer	7
Curd	10
Butter	30
Khoa	15

A longer shelf life allows for greater flexibility in transportation and marketing, which is crucial for rural areas with less developed infrastructure. Farmers should consider producing products like ghee that have fewer storage challenges.

Economic Impact of Value Addition on Rural Farmers

Table 6 evaluates the economic impact of value addition on rural farmers, focusing on income enhancement and employment generation. Farmers engaged in value-added processing experience a significant increase in income, with an average rise of 50-70% compared to those who sell raw milk. This increase is largely attributed to the higher market prices of processed products such as ghee, paneer, and butter, which have strong consumer demand. Additionally, value addition creates new employment opportunities in processing, packaging, and distribution, thereby contributing to rural job growth.

The table also highlights the multiplier effect of value addition on the rural economy, as the increased income allows farmers to invest more in their businesses, purchase better feed for their animals, and improve their overall standard of living. Employment generated through dairy processing further strengthens the local economy by providing additional

income sources for families involved in ancillary activities like packaging and transportation. The data clearly show that value addition has a transformative impact on rural livelihoods, not only by increasing direct income for farmers but also by stimulating broader economic growth in rural communities. This underscores the importance of promoting value-added dairy production as a strategy for rural development.

Table-6 Economic Impact of Value Addition on Rural Farmers

Impact Metric	Raw Milk Farmers	Value-Added Producers
Average Income Increase (%)	0	50
Employment Opportunities	Minimal	High

Conclusion

The study underscores the significant economic benefits of value addition in buffalo milk production for rural farmers in Mathura District, Uttar Pradesh. By transforming raw milk into products like ghee, paneer, butter, curd, and khoa, farmers experienced a substantial increase in income, with value-added products generating 50-70% higher earnings compared to raw milk sales. Ghee and paneer emerged as the most profitable due to their longer shelf life and high consumer demand, while curd and khoa, though profitable, required rapid market turnover. The data demonstrate that value addition can not only improve farmers' incomes but also stimulate rural employment and enhance the local economy.

However, the study also identified challenges such as limited access to modern processing technologies, inadequate market linkages, and a lack of technical training. Addressing these barriers through government policies and support systems can help maximize the benefits of value addition. Investments in dairy processing infrastructure, market access, and capacity building for small-scale farmers are essential to unlock the full potential of buffalo milk production. Overall, value addition represents a viable strategy for advancing the rural economy and improving the livelihoods of dairy farmers in India.

References

Subbanna Y B, Kumar S and Puttaraju S K M. 2021. Forecasting buffalo milk production in India: Time series approach. Buffalo Bulletin. **40**(2):335-343.

Khetra Y, Meena G S and Arora S. 2022. Buffalo milk and its products: Composition, nutrition, and benefits. In Chauhan M S and Selokar N (Eds), Biotechnological applications in buffalo research, Springer, Singapore, pp 67-89.

Kashyap D and Bhuyan S. 2023. Accessing value-added market through cooperatives: A case study of

Sitajakhala Dugdha Utpadak Samabai Samiti Ltd., India. Journal of Agribusiness in Developing and Emerging Economies. **13**(3):214-229.

Srivastava A K. 2015. Developing dairy enterprise for ensuring food and livelihood security of the poorest of poor. Indian Farming. **60**(10):17-20.

Muthukumar M, Kandeepan G, Pathak V, Rathod K S, Ambadkar R K, and Kulkarni V V. 2018. Carcass traits and value of meat and byproducts of buffalo. Indian Journal of Animal Sciences. **88**(3):326–330.

Journal of Rural Advancement
ISSN: 2347-2561 (P); 2583-6102 (E)
Website: jra.idtra.co.in

Instructions to the Authors

1. The following types of material will be considered for publication in the Journal (refer to guidelines for publication of research papers on the diseases not reported in India.

2.1 PAPERS ON ORIGINAL RESEARCH COMPLETED, not exceeding 3,500 words (approximately 10-12 typed pages), should be exclusive for the journal. They should present a connected picture of the investigation and should not be split into parts.

2.2 SHORT RESEARCH NOTES not more than 1,500 words (about 4-5 typed pages), which deals with (a) research results which are complete but do not warrant comprehensive treatment, and (b) descriptions of new material or improved techniques, with supporting data. Such notes require no headed sections. Summary (not more than 80-100 words) is to be provided at the end of the text.

2.3 CRITICAL RESEARCH REVIEW pointing out lacunae in research and suggesting possible lines of future work.

2.4 Contributors are requested to ensure that the research papers or notes submitted for publications have a direct bearing on animal production or open up new grounds for productive research. Basic types of papers and notes which relate to investigations in a narrow specialized branch of a discipline may not form an appropriate material for this journal.

3.1 The manuscript of the article starts with the title. It should be brief, specific and informative. It should be phrased to identify the content of the article and include the nature of the study, and technical approach, which is necessary for key-word indexing and information retrieval. Title should be as concise as possible, and include the theme involved in the research when applicable. Abbreviations should be avoided in the title.

3.2 In addition, a SHORT TITLE not exceeding 50 characters including space should be provided separately for running headlines.

3.3 The BY-LINE should contain, in addition to the names and initials of the authors, the place where the research work was conducted. Naming an author implies that the person named is aware of the research work reported and agrees with and accepts responsibility for any results or conclusions reported. The address of the institution should include the name of the department/Division, institution, city, country and pin code. Present address be given as a footnote. In case, a paper has several authors from different institutions, key the author to the address with superscript Arabic numerals and present the additional addresses as footnotes at the bottom of the page, e.g., Present address: Give designation, present address of all the authors and email of corresponding author. Designation, Division of... (1email of all authors); 2Designation, Division of...(if second author is from different division). Designation, Division of (if the author is from different place).

4. The ABSTRACT should be written in complete sentences and has not more than 250 words. It should contain a very brief account of the materials, methods, results, discussion and conclusion, so that the reader need not refer to the whole article except for details. It should not contain references to literature, illustrations and tables. The abstract should summarize pertinent results in concise way but understandable form. The abstract should start with a clear statement of the objectives of the research work and must conclude with one or two sentences that highlight important conclusions. "An abstract is defined as an abbreviated accurate representation of the contents of a document, preferably prepared by its author(s) for publication with it. Such abstracts are also useful in access [abstracting] publications and machine-readable databases".

4.1 Key-words: At the end of the abstract, list up to five to eight key words that best describe the nature of the research. Because major words in the title are not used in the subject index, appropriate words from the title (or synonyms) should be listed as key words.

5. Major headings are INTRODUCTION, MATERIALS AND METHODS, RESULTS AND DISCUSSION, and REFERENCES. Major headings of review papers or

papers from symposia may deviate from this standard format; however, all papers must contain an abstract, key words, and an introduction. Abbreviations should be avoided in headings.

5.1 The INTRODUCTORY part should be brief and limited to the statement of the importance of the study reported, problem or the objective(s) of the experiment. It may be a brief justification of the research work and specify the hypotheses to be tested. The review of literature should be pertinent to the problem. Objective of the study should be discussed in view of latest references. Authorities for the latin binomial of every organism are not used in the title or summary, and only on the first mention in the main body of the text. Gene names and loci should be italic, proteins should be roman. Virus nomenclature (and acronyms) should follow the guidelines of the International Committee on the Taxonomy of Viruses (ICTV). The current report is: van Regenmortel MHV, Fauquet CM, Bishop DHL (Eds) (2001) Virus Taxonomy: Seventh Report of the International Committee on Taxonomy Viruses. San Diego: Academic Press. Authors are also advised to check the ITCV website for the latest information. Biological nomenclature, as laid down in the International Code of Botanical Nomenclature, the International Code of Nomenclature of Bacteria, and the International Code of Zoological Nomenclature should be followed. www.sp2000.org and fishbase.org websites may be referred for nomenclature. Chemical nomenclature should follow the International Union of Pure and Applied Chemistry (IUPAC) definitive rules for nomenclature. No trade name should be used and Industrial products should be referred to by their chemical names (give ingredients in parentheses) at first mention. In the absence of a common name, use the full name or a defined abbreviation, in preference to a trade name. In case of anatomy and surgery articles please refer Nomina Anatomica, Nomina Histologica, Nomina Embryologica, Nomina Anatomica Veterinaria, and Nomina Anatomica cvium for nomenclature.

5.2 Relevant details should be given of the subject, MATERIALS AND METHODS, including experimental design and the techniques implied. Where the methods are well known, the citation of a standard work is adequate. All alterations of procedures must be explained. Experimental materials and statistical models should be clearly and fully described. Calculations and the validity of deductions made from them should be checked and validated by a statistician. When possible, results of similar experiments should be pooled statistically. Avoid to report a number of similar experiments separately. Units of measurement, symbols and standard abbreviations should conform to those recommended by the International Union of Bio-Chemistry (IUB) and the International Union of Pure and Applied Chemistry (IUPAC). Follow the Metric measurements and dosages should be expressed entirely in metric units (SI units). In exceptional circumstances, others may be used, provided they are consistent. Give the meaning of all symbols immediately after the equation in which they are first used. Equations should be numbered serially at the right-hand side in parentheses. In general, only equations explicitly referred to in the text need be numbered. The use of fractional powers instead of root signs is recommended. Powers of e are often more conveniently denoted by exp. In chemical formulae, valence of ions should be given as, e.g., Ca2+, not as Ca++. Isotope numbers should precede the symbols, e.g., 18O . The repeated writing of chemical formulae in the text is to be avoided where reasonably possible; in place of the name of the compound should be given in full. Exceptions may be made in the case of a very long name occurring very frequently or in the case of a compound being described as the end product of a gravimetric determination (e.g., phosphate as P2O5). Authors must certify that animals were cared for under guidelines comparable to those laid down by the Institutional Animal Ethics Committee.

5.3 The RESULTS AND DISCUSSION should preferably be combined to avoid repetition. Results should be presented in the tabular form and graphs when feasible but not both. The colour figures and plates, are printed in case only, when information would be lost if reproduced in black and white. Mean result with the relevant standard errors should be presented rather than detailed data. The data should be so arranged that the tables would fit in the normal layout of the page setup. Self- explanatory tables should be typed on separate sheets bearing appropriate titles. The tabular content should not exceed 20% of the total text. Any abbreviation used in a table must be defined in that table preferably as the footnote. Paginate the tables in respective series with the text. All tables should be cited in the text of the article. If an explanation is needed, use an abbreviation in the body of the table (e.g., DM) and explain clearly in footnotes what the abbreviation means. References to footnotes in a table are specified by superscript numbers, separately for each table. Superscript alphabets are used to designate statistical significance. Use a lower-case p to indicate probability values (i.e., $p<0.05$). In general, use numerals. When two numbers appear adjacent to each other, spell out the first (i.e., eight 1-day old chicks rather than 8 1-d old chicks). In a series using some numbers less than 10 and some more than 10 use numerals for all (i.e. 2 villages, 6 communities and 15

schools). Do not initiate a sentence with a numeral, it is better to spell it out or rearrange the sentence. Abbreviate the terms month (m), week (w), hour (h), minute (min) and second (sec) when used with a number in the text but spell them out when they are used alone. Do not use a hyphen to indicate inclusiveness (e.g., use 10 to 15 mg or wk 1 and 5 not 10-15 mg or wk 1-5). Use Arabic numerals with abbreviated units of measure: 12 g, 7 d, $10.00, 5% and numerical designations in the text: exp 2, group 4, etc. The images should be avoided if possible. The printing of figures needs additional payment of APC. If necessary, the author is required to submit high-resolution images, preferably with the initial submission but no later than revision stage. Electronic images (figures and schemes) must have a minimum resolution of 600 dpi for line drawings (black and white) and 300 dpi for colour or greyscale. Colour figures must be supplied in CMYK not RGB colours. Be ensured that the prepared electronic image files print at a legible size (with lettering of at least 2 mm). do not save the images as Portable Document Format (PDF). Please ensure that the figure is clearly labelled with its figure number. Original artwork: If you are unable to submit electronic artwork, please provide two sets of all figures as high-quality glossy prints at the size that they are to appear in print. One set should be unlabeled. The text content should elaborate on the tabular data, but not be repeated extensively within the text. Sufficient data, all with the index of variation attached, should be presented to allow the reader to interpret the results of the experiment. The discussion should interpret the results clearly and concisely in terms of proper mechanisms and should integrate literature results with the research findings to provide the reader with a wide base on which to accept or reject the hypotheses tested. Results and references to tables and figures already described in the RESULTS section therefore they should not be repeated in the DISCUSSION section.

5.4 The DISCUSSION should relate to the limitations or advantage of the author's experiments in comparison with the work of cited studies. Authors must obtain permission to reproduce any copyright material, and include an acknowledgment of the source in their article. They should be aware that the unreferenced use of the published and unpublished ideas, writing or illustrations of others, or submission of a complete paper under a new authorship in a different or the same language, comes under plagiarism. The Conclusion section should not be of more than one paragraph after the discussion. It should explain the implications of findings of this research on rural life in general terms. Abbreviations, acronyms, or citations should be avoided here. Though limited speculation is permitted, this section should also caution the reader against over extrapolation of results. For manuscripts with direct applications, this section will contain an interpretive summary. If results have no implications, this should also be stated.

5.5 A recent issue of The Journal of Rural Advances should be consulted for the methods of citation of References in the text as well as at the end of the article. Reference citations in the text are typed as follows: Martine (1971) or (Martine 1971); Black et al. (1974) or (black et al. 1974); Snedecor and Cochran (1994) or (Snedecor and Cochran 1994). Groups of references cited in a sentence in the text must be listed in chronological order as in the previous sentence.

5.6 REFERENCES lists should be given in alphabetic order. The reference list should be first sorted alphabetically by author(s) and secondly chronologically. For Journal Articles Author(s), year. Title. Journal title (full name and in italics) volume number (bold): page-page.

Bera I, Tyagi P K, Mir N A, Tyagi P K, Dev K, Sharma D and Mandal A B. 2019. Dietary supplementation of saponins to improve the quality and oxidative stability of broiler chicken meat. Journal of Food Science and Technology 56(3):2063-72.

For whole books Author(s), year. Title. (Ed.). Edition if any, Number of pages. Publisher, address.

Snedecor G W and Cochran W G. 1994. Statistical Methods, 6th edn, 491p. Iowa State University Press, Ames.

For chapters from books Author(s), year. Title. book title, editors (editors), Publisher, address. Page-page.

Clark J A and McArthur A J. 1994. Thermal Exchanges. Livestock Housing.1edn, pp. 97-122. (Eds) Wathes C M and Charles D R. CAB INTERNATIONAL, Wallingford, UK. For Symposium Devegowda G, Raju M V L N, Afzali N and Swamy H V L N. 1998. Mycotoxin picture world-wide: Novel solutions for their counteraction. Proceedings of 14th Alltech's Annual Symposium on Biotechnology in the Feed Industry. pp. 241-55. 5 May 1997. Bangalore.

Authors be ensured that all references in the text appear at the end of the paper and vice-versa, and that names and dates at the two places correspond.

6. All the articles are sent to referees for SCRUTINY and authors should meet criticism by improving the article.

7. Papers should be TYPE-WRITIEN on A4 paper, and double-spaced throughout (including references and tables) with a 5 cm margin at all the dimensions. Article (including illustrations) should be sent after a careful check-up of typographical errors.

8. Please upload the plagiarism checking report of your manuscript as supplementary file checked using popular software like iThenticate or Turnit-in.

9. Authors are requested consult The Council of Biology Editors Style Manual 7th edn, American Institute of Biological Sciences, Washington DC.

10. Proof-correction should be made in ink, in the margin. All queries marked in the article should be attempted and answered. Proofs are supplied for a check-up of the correctness of type-setting and facts. Excessive alteration may be charged to the authors. The proofs should be returned within 10 days.

11. Please click here to download article certificate and submit duly signed certificate (scanned copy only) may be attached online with article as supplementary file. Review Article Review articles on topical issues are invited from prominent scientists, peer-reviewed and published. Copyright Notice The copyright of the articles published in The Journal of Rural Advancement vests with the Journal, who has the right to enter into any agreement with any organization in India or abroad engaged in reprography, photocopying, storage and dissemination of information contained in these journals. The Journal of Rural Advancement has no objection in using the material, provided the information is being utilized for academic purpose but not for commercial use. Due credit line should be given to The Journal of Rural Advancement where information will be utilized.

Journal of Rural Advancement
ISSN: 2347-2561 (P); 2583-6102 (E)
Website: jra.idtra.co.in

Editorial Policies

Focus and Scope

Articles published in The Journal of Rural Advancement encompass a broad range of research topics related to rural life. The readership of the journal is global. The Journal of Rural Advancement is published both online and in print mode. The Journal of Rural Advancement includes articles on Agriculture, Animal husbandry, Rural culture and craft, Rural economics, Rural Education, Rural sociology, Rural health and care, Rural welfare, and Science and technology.

Section Policies

Sections	Checked Open Submissions	Checked Indexed	Checked Peer Reviewed
Review Article	Yes	Yes	Yes
Articles	Yes	Yes	Yes
Short-Communication	Yes	Yes	Yes
Special Feature	Yes	Yes	Yes
Invited Review	Yes	Yes	Yes
Graphical Abstract	Yes	Yes	Yes

Peer Review Process

Each paper submitted for publication is sent to independent referees for peer review. They are selected on expertise in one or more areas of paper, no conflicts of interest, ability to think clearly and logically, ability to write a good critique, accurate, reliable in returning reviews, and ability to do the review in the allotted time frame.

Referees are expected to respond to the editor's request for advice within a limited period (15 days), which is clearly stated by the editor. A comment sheet is also provided to him for seeking his advice on all aspects of the article. He is advised to return the script immediately without comments if he can't attend to a manuscript within this period so that the editor can send it to another referee without further delay.

Comments from the referee are received by the editor

If the reports of both referees disagree concerning the suitability of the paper for publication, the advice of a third referee is sought.

Comments received from both referees: The article is sent to the corresponding author through email only for modification along with comments from referees. At this stage, the editor completes the preliminary editing. It includes checking in-house style (presentation of title, affiliations, present address, abstract, scientific nomenclature, text, and reference presentation).

Comments not received from both referees: If comments are not received from both referees, a reminder is sent to them.

Reminder to referee from whom the comments are not received: If comments are not received from one referee, a reminder is sent to him.

Comments not received from one referee: A reminder is sent to the referee who is not responding.

A reminder is sent to the referee who is not responding to the editor's request. However, if his reply is not coming in a stipulated time, the editor takes suitable action based on comments received from one referee.

Rejection of the article based on comments of referees: The article is rejected based on the comments of referees/referees. The corresponding author is informed accordingly.

The corresponding author does not agree with the adverse comments of the referee or in the meantime, suitable comments are received from the second referee. Editor's decision independently or in consultation with the editorial board is final and not challengeable elsewhere. The editor may ask the first referee or a third referee to go through the comments of the first referee and reply to communicating author to comments of the referee and /or contradictory comments of the second referee.

Favorable comments are received from the second referee after the modification: Input of the second referee is incorporated in the modified article. The corresponding author is contacted if the referee asks for the inclusion of more information in the form of a table, figs, or references.

Adverse comments received from the second referee at any stage: The corresponding author is informed of the adverse comments from the second referee and action is stalled (even at the proof stage) till the final decision.

The editor's decision independently or in consultation with the editorial board is final. Acceptance of the article is sent to the corresponding author.

Publication Frequency

The Journal of Rural Advancement is a Biannual Journal. However, the article will be published online immediately after its acceptance.

Publication Language

The language of the journal is English.

Open Access Policy

This journal provides immediate open access to its content on the principle that making research freely available to the public supports a greater global exchange of knowledge.

Journal of Rural Advancement
ISSN: 2347-2561 (P); 2583-6102 (E)
Website: jra.idtra.co.in

ARTICLE CERTIFICATE

Article. No.
Article Entitled:
.....
.....
Title changed to:
.....
.....
Authors (names):
.....
.....
It is certified that:

1. The article has been seen by all the authors (signatures given below) and are satisfied with its form and content.
2. The sequence of names of the authors in the by-line is as per their relative contribution to this work, giving due credit to all scientists who made a notable contribution to it.
3. The address of the organization where the research was conducted is given in the by-line (changes of authors address is provided in the footnote).
4. The experiment(s) was conducted during 20..... The article is submitted to the journal for publication soon after completion of the experiment(s).
5. The article is exclusive to this journal, and the results reported here have not been sent (and not be sent during its consideration by this journal) for publication in any other journal.
6. The article has not been rejected for publication in any other journal, been rejected in .... whose comments are attached (such a rejection does not disqualify the article for consideration in this journal).
7. The work is based/ not based on a part/ complete M.Sc/Ph.D. thesis of the first author, who submitted the thesis in 20..... to the ..... (university, place).
8. Original and genuine data and facts are presented in the article.
9. I/ we agree to abide by the objective comments of referees and do agree to modify the article into a short note as per the recommendation, for publication in The Journal of Rural Advancement.
10. If published in The Journal of Rural Advancement, the copyright of this article would vest with the Journal and will have the right to enter into any agreement with any organization in India or abroad engaged in reprography, photocopying, storage, and dissemination of information contained in it, and neither we nor our legal heirs will have any royalty claims.
11. Article processing fee deposited in the bank (Account holder: Vraddhi Organic Agro India Private Limited, Account No. 22230200000799, The Federal bank Limited, Vrindavan, Branch code: 2223, IFSC: FDRL0002223, SWIFT CODE: FDRLINBBIBD).
INR/$ ….....
Date ….....
UPI ….....


The information furnished above is true, correct, and complete to the best of my/our knowledge and belief.

Sl. No.	Signature	Date	Name in full and designation	Present official address

Note: Extra sheet(s) may be attached if needed.

Journal of Rural Advancement

[ISSN 2347-2561 (P); 2583-6102(E)]



Vol.12 Issue 2 October 2024

Official Journal of
Institute for Development of
Technology for Rural
Advancement

IDTRA